

implants

international magazine of oral implantology

4²⁰¹⁵

| **research**

Incidence of postoperative infections in dental procedures

| **case report**

Abutment fracture in a bridge supported by natural teeth and implants

| **industry report**

Complete reconstruction for a patient with chronic tooth decay

oam113

Where innovation comes to life

Nobel Biocare Global Symposium

June 23–26, 2016 – New York



A tradition unlike any other

The 150 exceptional speakers, program and venue make the Nobel Biocare Global Symposium a must-attend event for dental professionals worldwide. Held at the Waldorf Astoria in New York, it's a unique opportunity to experience how innovation can come to life in your daily work.

Design your own learning experience

Choose from numerous podium lectures, forums, master classes and hands-on sessions – with some of the best speakers and presenters in the world. The exciting program will be divided into three main themes: refining treatment, digital dentistry and challenging clinical situations. Don't miss this opportunity. Sign up now and we'll see you in New York!

**Register
now**

nobelbiocare.com/global-symposium-2016



Implantology & Education



Dr Rolf Vollmer

As 2015, an eventful year of implantology, is coming to a close, a review on the DGZI's activities shows its great dedication to implantology. The German Association for Dental Implantology (DGZI) has been active both in Germany and abroad and its cooperative partners, its various resources and concepts will make sure that the DGZI continues to be a significant influence on implantological education around the world.

A special concern of the DGZI's activities this year was the promotion of dental technologies with regard to a successful implant therapy. This year's 45th International Annual Congress, which was visited by more than 350 participants, was therefore dedicated to the interface between dental technology and implantology. International attendees were able to participate in all specialist discussions and to present their own key topics with the help of interpreters and special international podiums.

We already look forward to welcoming the participants at next year's workshops and our International Annual Congress, which will take place in Munich from 30 September to 1 October 2016. Thus, you can expect a special Octoberfest Congress on the last weekend of the "Wiesn".

In addition, I would like to thank you all for the provision of various specialist articles this year and ask you to continue your expert work for **implants international magazine of oral implantology**. Like in the previous years, all articles will be reviewed by our scientific committee.

Moreover, **implants international magazine of oral implantology** is distinguished from other publications by presenting controversial topics. A lively discussion and even occasional topic-related disputes among colleagues form an important part of striving for the best-possible therapeutic approaches in implantology.


The members of the DGZI board hope you will enjoy reading this year's final edition of **implants international magazine of oral implantology** and wish you a happy and peaceful Christmas as well as a good start to 2016.

Dr Rolf Vollmer


First Vice-President and Treasurer of the German Association of Dental Implantology



page 6



page 14



page 26

| editorial

03 Implantology & Education

| Dr Rolf Vollmer

| research

06 Incidence of postoperative infections in dental procedures

| Prof. Mauro Labanca

| case report

14 Abutment fracture in a bridge supported by natural teeth and implants

| Dr Gregory-George Zafiroopoulos *et al.*

20 Minimally invasive implant placement without the use of biomaterials — The bone expansion technique

| Dr Gilles Chaumanet

26 CAD/CAM custom-milled titanium bar for rehabilitation of an atrophic upper jaw

| Dr Richard Marcelat

| industry

32 Complete reconstruction for a patient with chronic tooth decay — The damage undone

| Dr Ara Nazarian

38 Digital precision for all indications

| Nobel Biocare

| events

42 “Our members profit from a wealth of international experience” — 45th Annual International Congress of the DGZI

| DGZI

46 50 Years of clinical osseointegration

| Georg Isbaner

| news

40 Manufacturer News

48 News

| about the publisher

50 | imprint



Cover image courtesy of DENTSPLY IH GmbH
www.dentsplyimplants.de
Original Background: © Vik Y
Artwork by Sarah Fuhrmann, OEMUS MEDIA AG.



page 32



page 42



page 46

Restoring happiness

Patients rely on you in order to eat, speak, and smile with confidence. It can be said, you are actually restoring quality of life and happiness.

To succeed, you need technology that is well founded and documented in science. That is why we only deliver premium solutions for all phases of implant therapy, which have been extensively tested and clinically proven to provide lifelong function and esthetics.

Moreover, with an open-minded approach, we partner with our customers and offer services that go beyond products, such as educational opportunities and practice development programs so you can rest assured that you have the support you need.

Reliable solutions and partnership for restoring happiness—because it matters.

Comprehensive solutions for all phases of implant dentistry




Professional and practice development

STEPPS™



Digital planning

SIMPLANT®



Regenerative solutions

SYMBIOS®



Implants

ANKYLOS®

**ASTRA TECH
IMPLANT SYSTEM**

XiVE®



Restorations

ATLANTIS®

Incidence of postoperative infections in dental procedures

Author_Prof. Mauro Labanca, Italy

The risk of post-surgical infections in dental surgery and the management of the same has been widely studied and referenced in medical literature.¹

Actually, it is known that in order for any surgical wound to heal properly and in a predictable manner, two conditions that I would define as "milestones" must be met: the wound should be protected from any trauma and prevent superinfection of the same. These two conditions can hardly be met in the oral cavity. It is well known that the oral cavity, which is the first section of the digestive system, is an intrinsically contaminated environment and the risk of infection during intra-oral surgery is increased compared to other types of surgeries and comparable to surgery on the intestine. In fact, it is practically impossible to en-

sure an aseptic environment due to the large number of microorganisms present in the oral cavity; as we all know, bacteria, fungi and protozoa live in the soft tissues creating a biofilm. The life cycle of the biofilm depends on the attack, the colonisation and the proliferation of these micro-organisms.

Common bacterial flora in the oral cavity is variable and consists of aerobic and anaerobic bacteria with pathogenic potential.² Temporary reduction of the amount of such bacteria may reduce the risk of post-surgical infection.³

Therefore, before performing a surgical procedure, it is essential to consider that the wound is never sterile and when subjected to an infection, the latter is due to perioral skin microflora.

Moreover, surgical wounds caused by dental surgery are continuously subjected to trauma: mastication, dental prosthesis, movement of the tongue or perioral muscles. In fact, this involuntary and persistent trauma cannot be eliminated in any way and, obviously, affects significantly the wound healing time.

Speaking with my youngest colleagues as well, they usually pay a great deal of attention to the operating sequence ignoring the key factor that conditions the outcome of the intervention: the proper healing of the wound. In fact, if there is a superinfection of the wound with consequent dehiscence of the flap, the intervention itself and/or the grafted material are likely to turn into failure or unsuccessful operation. Therefore, it is critical to set wound healing as the main goal

Fig. 1_Severe facial infection after mandibular third molar extraction.



Implant planning made easy

Planmeca PlanScan®

Planmeca ProMax® 3D

3x
3D

Planmeca Romexis® software offers a completely integrated and digital workflow for modern implantology. From intraoral scanning to easy prosthetic design and realistic implant libraries, the most sophisticated implant planning tools are just a few clicks away.

- All the scanned and design data for prosthetic works is immediately available and can be mapped with the patient's CBCT data
- Use crown library or patient-specific crown from CAD system
- Verify the implant plan with verification tool
- Order surgical guide directly from the software
- Share data easily with partners through **Planmeca Romexis® Cloud** image transfer service



Find more info and your local dealer

www.planmeca.com

PLANMECA



Planmeca Oy Asentajankatu 6, 00880 Helsinki, Finland. Tel. +358 20 7795 500, fax +358 20 7795 555, sales@planmeca.com



Fig. 2

Fig. 2_ Adequately prepared surgical table.

and make sure the wound is protected against superinfection and trauma, although it is located in a dynamic and contaminated environment.

But how can a post-surgical infection be prevented?

Based on medical literature and my extensive experience as an ER surgeon and dental surgeon, I think that there are different parameters to take into consideration when performing a dental surgery: the experience of the surgeon, the duration of the surgery itself, the concomitant risk factors, the aseptic conditions of the operating field and the careful selection of the materials used.

It is also essential to keep in mind that oral surgery is not just about implants or various regenerative techniques. Even a seemingly common avulsion can be fully considered a surgical intervention, hence subject to infection with more or less serious side effects for the patient (Fig.1).

Fig. 3_ Incorrect preparation of a surgical table.




Fig. 3

We will now review the above risk factors for an in-depth examination of every single situation.

Experience

Experience proves to be the most important factor in a successful outcome of a surgical intervention. It has been reported that the risk of infection in the case of less experienced surgeons is four times higher compared to that of more experienced surgeons.⁴ Nevertheless, experience is definitely not a parameter that can be changed (unless by aging and through hard work!) but it is necessary to take note of it, and then young colleagues who face surgery should pay more attention to their work aware of this aspect.

Duration

In defining the duration of the intervention, there are two factors that must be considered: the duration, in a relative sense and in an absolute sense. The absolute value indicates the time required for the execution of the surgical procedure in optimum conditions by a surgeon with adequate experience. Virtually, the right execution time, with no rush but also without unnecessary expenditure of time. On the other hand, when lack of experience or insecurity lead to extended duration, we talk about relative value: basically, it is the time actually spent but that could have been reduced. It has been reported that a duration of the intervention below one hour poses a risk of superinfection of 1.3 %, while such risk is increased to 4% if the intervention lasts for about three hours. Every additional hour doubles the risk of superinfection⁵. Once again, it should be mentioned that these values refer to the correct duration of the intervention. To clear this up, if an intervention executed correctly lasts one hour, the risk is 1.3 %, if it lasts three hours, the risk rises to 4 %. But if the relative intervention can be done in one hour, but it takes three hours due to surgeon's lack of experience, the risk of infection increases considerably beyond the above said 1.3 %.

Systemic factors

There are systemic factors that promote superinfection of the blood clot including uncompensated diabetes (which also prolongs healing time much more than usual), autoimmune and systemic disorders, and smoking.^{6,7,8} Concomitant use of drugs should also be carefully considered to avoid that some of them could heavily interfere with the healing process (just think of the bisphosphonates, a problem that is more and more present in our clinical activity). The age of the patient should also be carefully evaluated; during the avulsion procedure of a third molar, the risk of post-surgery complications is of 10% in twenty-years-of-age patients and 30% in a 40-year-old patient.⁹ Actually, age involves very often the use of medicines and impaired

MORE BONE Where it Matters Most...



Find out more about the new V3 Implant at: www.V3-implant.com

V3 By **MIS**