DENTAL TRIBUNE

—The World's Dental Newspaper · Middle East & Africa Edition ——

Published in Dubai April 2011 No. 4 Vol. 9



News & Opinions
Events in the Middle East

Page 5



Media CME Diode laser Surface

▶ Page **6**



Dental CafeGet eco-friendly Bra!

• Page 11

WHO takes on global influenza threat

Daniel Zimmermann

Oberuzwil, **Switzerland** /LEIPZIG, Germany: Member of a working group set up by the World Health Organisation have agreed upon an international framework to enhance preparedness for influenza pandemics that threaten public health worldwide. The agreement, which is expected to provide clear legal regimes and responsibilities for all stakeholders involved in the prevention and management of pandemics, is the result of more than three years of negotiations which started back in 2007 during the height of the bird-flu influenza virus in Southeast Asia. It is expected to be ratified during the World Health Assembly in Geneva in

According to a joint statement, one of the key elements of the agreement will be an improved cooperation and exchange of information between key players such as the WHO, national laboratories and pharmaceutical manufacturers. Access

Switzerland in May.

to live-saving vaccines and other resources for low-income countries which often cannot produce or afford required anti-viral medication for their population is also supposed to be improved.

"This agreement promotes global health security and solidarity in pandemic times, " said Ambassador Bente Angell-Hansen, who also chaired the working group. "It also reflects a unique partnership with industry and contains concrete meas-

ures of cooperation with both industry and civil society."

Owing to increasing global transportation, locally active influenza viruses exhibit an increasing potential to become global pandemics that risks the life of many including medical and dental professionals. According to latest estimates of the WHO, the H1N1-virus or swine flu that first occurred in Mexico in early 2009 has killed almost 20,000 people worldwide.



Periodontal treatment no harm to newborns



NEWYORK, USA/LEIPZIG, Germany: Pregnant women with gum disease may undergo non-surgical periodontal treatment without fear of consequences for their baby's health. In a large trial involving 400 infants between the ages of two and three from different paediatric clinics in the US, dental clinicians found that treating periodontitis during pregnancy did not affect the children's cognitive, motor or language capabilities later in life.

In the study, clinicians from universities in Minnesota, Ken-

tucky, Mississippi and New York compared development data of children born to women who were treated for gum disease before and after their delivery. However, the results between the control and experimental groups only differed slightly. Higher motor and cognitive scores were observed in the children of women who saw an improvement in their periodontal health

Earlier studies indicated that paternal periodontal treatment

 \rightarrow DT page 3


PROMEDICA

Highest quality made in Germany

- high quality glass ionomer cements
- ► first class composites
- innovative compomers
- modern bonding systems
- materials for long-term prophylaxis
- temporary solutions
- bleaching products...

All our products convince by

- excellent physical properties
- perfect aesthetical results



Temporary crown and bridge material



Light-curing nano-ceram composite

- universal for all cavity classes
- comfortable handling, easy modellation
- highly esthetic and biocompatible





Technology

• exact match of translucency and shade to the packable Composan bio-esthetic

particular fracture and wear resistance now available in 6 attractive shades Resin-reinforced glass ionomer luting cement

- strong adhesion, very low film thickness
- especially suited for zirconia-based pieces



PROMEDICA Dental Material GmbH

Tel. + 49 43 21 / 5 41 73 · Fax + 49 43 21 / 5 19 08 Internet: http://www.promedica.de · eMail: info@promedica.de

Production manager

Dubai Health Authority signs agreement with Green Crescent **Insurance Company**



The Dubai Health Authority signed a service level agreement with Green Crescent Insurance Company, today, at the DHA headquarters making Green Crescent the first insurance company in UAE to have direct billing access to the DHAPublicHospitals and PrimaryHealthCareCenters.

The agreement was signed by His Excellency Khalid Al Sheikh

Editorial Board

Prof. Abdullah Al-Shammary, Restorative Dentistry, KSA Prof. Hussain F. Al Huwaizi, Endodontics, Iraq Prof. Samar Burgan, Oral Medicine, Jordan Dr. Abdel Salam Al Askary, Implantology, Egypt

Dr. Talal Al-Harbi, Orthodontist, Qatar Dr. Mohammed H. Al Jishi, Bahrain Dr. Lara Bakaeen, Prosthodontist, Jordan Dr. Aisha Sultan, Periodontist, UAE Dr. Kamal Balaghi Mobin Aesthetics, Iran

Published by Education Zone

 $in\ licence\ of\ Dental\ Tribune\ International\ GmbH$

© 2011, Dental Tribune International GmbH. All rights reserved.

Dental Tribune makes every effort to report clinical information and manufacturer's product news accurately, but cannot assume responsibility for the validity of product claims, or for typographical errors. The publishers also do not assume responsibility for product names or claims, or statements made by advertisers. Opinions expressed by authors are their own and may not reflect those of Dental Tribune International.

President/CEO Yasir Allawi y.allawi@dental-tribune.ae

Director mCME: Dr. D. Mollova

Marketing manager Khawla Najib


info@cappmea.com

khawla@dental-tribune.ae

dentalme@dental-tribune.ae

PO Box 214592, Dubai, UAE, Tel + 971 4 391 0257 Fax + 971 + 366 + 4512 www.dental-tribune.com

AD



NICOLAS ASP UNIVERSITY COLLEGE Dubai - Abu Dhabi

First Postgraduate Dental Institution in UAE

All Programs Accredited by UAE Ministry of Higher Education & Scientific Research

Register NOW...

MASTER DEGREE IN ORTHODONTICS & MALMO UNIVERSITY SPECIALIST DEGREE

Full and Part Time



SPECIALITY CERTIFICATE IN ORTHODONTICS

Full and Part Time

Swedish Recognition in Progress

Full and Part Time

MASTER DEGREE IN PEDIATRIC DENTISTRY

SPECIALITY CERTIFICATE IN PEDIATRIC DENTISTRY Full and Part Time

MASTER DEGREE IN ENDODONTICS

Swedish Recognition in Progress Full and Part Time

SPECIALITY CERTIFICATE IN ENDODONTICS

Full and Part Time

DIPLOMA IN ADVANCED EDUCATION IN **GENERAL DENTISTRY**

Fulfill Experience Requirements Full Time

HIGH DIPLOMA IN ORAL IMPLANTOLOGY

Modular option Available Full and Part Time









Abu Dhabi Headquarters

Admission Office: Al Nadi Al Seyahi Sheikh Omar Bin Zayed Al Nahyan

Dubai Branch

Dubai Healthcare City Ibn Sina Building (No. 27) Block D, 3rd floor, Office 302 Telephone: +971 4 3624787

Fax: +971 4 362 4793 P.O. Box: 53382, Dubai, UAE Website: www.dubaipostgraduate.com Email: info@dubaipostgraduate.com

Mubarak, Deputy Director General of the DHA and Carl J. Sardegna CEO of Green Crescent Insurance Company.

The agreement provides 110,000 Green Crescent insurance card holders with access to all DHA hospitals and health cen-

Saudi Arabia seeks designers for medical cities

The Saud health ministry has approached seven expert international firms to design the five medical cities ordered by King Abdullah, the Saudi Gazette has reported. In addition to the medical cities, which will be operational within five years, the number of beds at the ministry's hospitals will be increased by 70% in the next four years, from 34,200 to 54,000 beds at more than 349 hospitals and 2,750 primary care centres.

Qatar set for new oral health strategy

Qatar's Supreme Council of Health (SCH) is creating a national oral health strategy. The strategy is expected to be formulated after completing a nationwide survey on oral health among children and teenagers. Dental caries and periodontal diseases are the most prevalent among schoolchildren and teenagers in the Gulf state. The study, which will begin in the last quarter of this year, will help improve the existing oral health services and programmes in the country as well as assist to monitor the trends of oral and dental diseases in the country, SCH said.

Opening of Al Barsha Medical Centre

Qadhi Saeed Al Murooshid, Director General of the Dubai Health Authority (DHA) laid the foundation stone for the state-ofthe-art Al Barsha Health Centre. The centre, which is located in Al Barsha 2, has a built up area of 70,000 sq ft, will be constructed with an investment of Dhs50m. The project will be completed by mid 2012 and will serve a population of approximately 60,000 people living in the vicinity.

Bugs threaten health of orthodontic patients

Daniel Zimmermann

DTI

LONDON, UK/LEIPZIG, Germany: Orthodontic retainers are a potential source of harmful microbes if not properly cleaned, scientists in the UK have warned. In a series of tests conducted at the UCL Eastman Dental Institute in London at least 50 per cent of all tested retainers contained species of Candida and Staphylococcus micro-organisms, including MRSA, a multidrug-resistant bacterium that can be fa-

tal to patients with a compromised immune system.

The Candida yeast, found universally on human skin and other areas, can also cause infections. Amongst other conditions, it has been associated with oral candidiasis, a condition often related to ill-fitting dentures.

Both species do not normally occur in the oral cavity.

The researchers said that the high number of harmful bacteria found in retainers is most likely the result of poor cleaning, allowing microbes to build up a re-

powder chamber.

settings.

Uniquely simple to use.


Incredibly easy to operate.

Touch 'n' flow: Highly sensitive

3-touch panel for easy choice of

sistant biofilm and spread to other areas of the oral cavity such as interior cheeks and tongue. The potential for transmission is also high, as retainers are frequently removed and replaced in the mouth by the person who uses it, they added.

They recommend wearers wash their hands thoroughly before and after inserting their retainers. Proper dental hygiene through tooth brushing and the use of mouthwash also helps to keep harmful bacteria from entering the mouth.



For more information > welcome@ems-ch.com

AD

Iraqi doctors meet in Sharjah

WAM Sharjah, Sheikh Mohammed bin Saqr Al Qasimi, Assistant Undersecretary of Ministry of Health, opened the 6th conference of the International Iraqi Medical Association (IIMA) in collaboration with Sharjah Medical District under the auspices of UAE Health Minister Dr. Hanif Hassan and participation of Royal College of Surgeons in Ireland..


Addressing the medical gathering of 700 doctors of diverse specialisations, Sheikh Saqr said the three-day conference provides an ideal platform for sharing latest medical developments, knowledge and experiences.

About 75 speakers delivered 93 presentations through 18 panel of discussions. IIMA is a non-government organization (NGO) registered with the United Nations and seeks to raising awareness and increasing international exposure of Iraqi physicians and their contributions and services to the global medical profession, working and cooperating with prominent international scientific figures in the medical and healthcare industry.



may be linked to different medical problems including low birth weight, preterm birth and long-term development delays, as bacteria released during treatment may enter the mother's bloodstream and harm the baby. According to research, pregnant women are prone to gingival bleeding, which is caused by a hormonal imbalance that encourages the growth of certain oral bacteria.

If the new data is verified, pregnant women throughout the US could have their gum conditions treated, confident in the knowledge that it will not have a clinically significant effect on their child's development, the researchers said. A spokesperson of the American Academy of Periodontology said that although the data remains inconclusive, the organisation generally recommends women to maintain their periodontal health during pregnancy.



Industry veteran honoured with first DT Award 2011 nominations are now open at Dental Tribune website

Yvonne Bachmann

 $\overline{\mathrm{DTI}}$

COLOGNE & LEIPZIG, Germany: Robert Gottlander from Sweden was announced the first winner of the Dental Tribune Industry Leadership Award, an honour for outstanding achievements in dental education and innovation. Gottlander received the award from DTI CEO Torsten Oemus during a recent award reception at IDS Cologne.

Gottlander attended the School of Dentistry at the University of Gothenburg. He started his career at Nobelpharma in 1984, which was later consolidated into Nobel Biocare.

During his time as executive vice president for education and products, from 1986 to 1987, Gottlander was responsible for the internationally acclaimed educational training programme, Nobel World Tour.

Being awarded annually, the Dental Tribune International's Global Dental Tribune Awards aims to recognise outstanding individuals, teams and practices that have an active interest in continuing professional development and staying at the top of the profession.

The audience consists of over 650,000 dental professionals, all readers of the Dental Tribune newspaper, which the net-

work is publishing in more than 25 languages.

Mr Oemus said that 15 categories are now open for nominations including Lifetime Achievement, Innovation in Dentistry or Dental Educator of the Year Award at the Dental Tribune Website (www. dentaltribune.com/awards.) The winners will be announced at this year's Greater New York Dental Meeting in November.



Torsten Oemus (right) handing over the first Dental Tribune Award to Robert Gottlander. (DTI/Photo Yvonne Bachmann, DTI)

Scientists declare CUS an autoimmune disease



NEW YORK, USA/LEIPZIG, Germany: Patients that suffer from a very rare condition affecting the oral mucosa may soon get relief from new research conducted in the US. In a recent study, scientists from the Tufts University near Boston claim to have found evidence that the socalled Chronic Ulcerative Stomatitis (CUS), characterised by recurring painful ulcers, is mainly caused by an autoimmune response of the body that destroys the binding of cells inside the surface tissue layer of the mouth.

According to the scientist, only a dozen cases of CUS have been reported worldwide since the condition was first clinically identified in 1989 but more patients could be affected due to the extensive testing procedure and low awareness among dental clinicians.


They said although prior it was known that affected patients had specific autoantibodies, researchers were not able to determine how much these actually contributed to the condition.


With help of the new findings, CUS could now be classified as an autoimmune disease in order to allow better management of the symptoms.


Due to its unique resistance to standard medication like corticosteroids, successful treatment of CUS has been achieved only in some cases through hydroxychloroquine, a prescription drug primarely used to prevent malaria as well as to treat rheumatoid arthritis or lupus. By better understanding the mecha-

nisms linking the autoimmune response to ulcerative sores, new treatment approaches could be developed to manage the condition, the scientists said.

So far, CUS has been found primarily in middle-aged Caucasian woman. It can only be diagnosed by surgical biopsy using immunofluorescence microscopic examination in an outside lab. In normal clinical settings it can be taken for oral erosive lichen planus, another more common chronic condition affecting mucosal surfaces and also considered to be an autoimmune disease.







CONGRESSES

May 1, 2011 - May 5, 2011 -**Dubai**, UAE

Dubai Oral & Maxillofacial Surgery Congress Web: www.iir.dubaiomfs.org Email: meetings@imedex.com Venue: Westin Dubai Mina Seyahi

May 6, 2011 - May 7, 2011 -Beirut, Lebanon

ITI Congress Middle East Web: www.iti.org/congressmiddleeast Email: events@iticenter.ch

May 12, 2011 - May 13, 2011 -**Dubai**, UAE

5th CAD/CAM & Computerized Dentistry International Confer-

Web: www.cappmea.com Email: info@cappmea.com

DT Group extends to platform

Daniel Zimmermann

Members of the Dental Tribune International Publishing Group (DTI) met in Cologne recently for the 7th Annual Dental Tribune International Publishers Meeting. The gathering, which is traditionally held prior to IDS, saw new licence partners from Slovenia, the Netherlands and the Czech Republic joining the Group. The largest global dental publisher's network now comprises 28 partners, including Russia, China and India, to name a few.

New features of DTI's online portfolio were also revealed in Cologne. According to Publisher and CEO Torsten Oemus, users of the website www.dental-tribune. com will now be able to post and search jobs and classifieds worldwide and in their respective local markets. He also announced a free app featuring a selection of news articles and videos from the DTI network and in different languages for Apple's iPhone and iPad.

Based in Hong Kong, New Yorkand Leipzig in Germany, DTI currentlypublishes over 100 newspapers and magazines in 90 countries worldwide. Their offering is extended by online education realised by the Dental Tribune Study Club, DI

Events in the Middle East

Jun 2, 2011 - Jun 4, 2011 - Istanbul, Turkey

European Academy of Esthetic Dentistry Spring Meeting 2011 "Controversies in Esthetic Dentistry E-mail:

ersoytabaklar@plazatur.com Venue: Swissotel The Bosphorus

TRADE SHOW

Jun 11, 2011 - Jun 14, 2011 -Tehran, Iran 14th IranMed 2011 E-mail: info@iranfair.com Web: www.iranfair.com //

www.iranmedonline.com

Jun 19, 2011 - Jun 23, 2011 - Istanbul

87th Congress of the European **Orthodontic Society** E-mail: cnidus@cnidus.com Website: www.cnidus.com Venue: Istanbul Convention & Exhibition Center (ICEC)

Oct 8, 2011 - Oct 9, 2011 - Istanbul, Turkey

Dental Istanbul E-mail: info@dentalistanbul.com// bilgi@vestiyer.com.tr Website:

www.dentalistanbul.com// www.vestiyer.com.tr

AD






Implant System


For more than 30 years, Dentoflex has been creating healthy and attractive smiles in Brazil, and is now growing all over the world.


Dentoflex goes beyond


Dentoflex has been operating in the dental market for over three decades and doesn't stop reinventing itself. The company is moving in the right direction by aiming to develop on a daily basis, invest in the latest technology and expand into new horizons. Dentoflex has an ambition to break down and overcome any barriers and form strong relationships and alliances in countries worldwide.

Treatment Surface









The Dentoflex's surface treatments includes aluminum oxide blasting, followed by acid passivation which combines texture and perfect porosity for an excellent biological effect, and as a result giving conditions for osseointegration and clinical success. The Dentoflex implants offer a more rougher surface of up to 300 times greater than a machined surface, stabalising bone-implant contact by up to six times.











BRANCH OFICCE: JAFZA VIEW - LOB 19 OFFICE 1301 - P.O. BOX 261901 - DUBAI - U.A.E TEL: +971 (0) 48855404 EXT: 304/305 - FAX:+971 (0) 48855405 MOB: +971 (0) 559250844 comex@dentoflex.com.br www.dentoflex.com.br



Diode laser surface decontamination in periodontitis therapy

(mCME articles in Dental Tribune (always page 6) has been approved by HAAD as having educational content acceptable for (Category 1) CME credit hours. Term of approval covers issues published within one year from the distribution date (September, 2010). This (Volume/Issue) has been approved by HAAD for 2 CME credit hours.

_We don't always have the opportunity to provide long-term dental treatment for patients with a profound marginal parodontopathy who have undergone resective surgical therapy, at times including reconstructive work. Correspondingly, there is only a limited amount of literature available due to the aforementioned fact. The number of published studies/other publications is even more limited as regards new therapy concepts or adjuvant treatments to complement a proven therapy regimen. In 1995, the first diode laser (wavelength 810 nm) was presented at IDS in Cologne. This device-initially as a prototypehad been used within the scope of a test phase since 1994. At the end of 1994 patients were treated with this "new" laser wavelength for the first time, which had not been used in dentistry up until that time. The Freiburg laser work group led by Krekeler and Bach, who were the first ones to deal with the integration of diode laser light in dentistry, noticed the considerable advantages of this new technology.

High-performance lasers emit monochromatic coherent light at a wavelength of 810 nm. This light is absorbed particularly well by dark surfaces. Thus the injection laser (= diode laser) is ideally suited to perform cuts, as are common in dental surgery, as well as for the removal of benign tumors in the oral cavity, for exposing implants and for use in mucogingival surgery. This excellent cutting performance of the diode laser can be attributed to the exceptional absorption of the laser light by the hemoglobin in the tissue. Aside from an application in softtissue surgery, the diode laser is also used for decontaminating surfaces that are colonized by germs (on implants and teeth). It was proven in these applications that especially a gram-negative, anaerobic germ spectrum is sufficiently damaged by the laser light.

The following paper describes-by means of three selected patient cases-our "Freiburg" experience of incorporating laser light decontamination in the therapy of marginal parodontopathies.

_Material and methodology

We are presenting treatment results for three patients who received dental treatment over a period of 15 years (12/94-04/10). Initially, these three patients suffered from a profound parodontopathy with inadequate degeneration of supportive tissue. The course of treatment for these three patients proceeded according to the following regimen:

1. Initial therapy (12-1994 through 01-1995)

- Motivation and instruction of the patient
- Cleaning and polishing
- Application of disinfecting
- 2. Resective phase (01-1995 and/or 02-1995)
- Creation of a mucoperiosteal
- Removal of granulation tis-
- Decontamination with diode laser light (p=1.0 Watt; tmax=20 sec)
- Apical shifting of soft tissue 3. Reconstructive phase (01-1995 and/or 02-1995)

Fig. 6_Orthopantomogram after insertion of 3 short implants ("shorties") in the atrophied left half of the mandible.

Fig. 7_6-year follow-up in 2001. Fig. 8-11_Clinical findings in 2009, shortly prior to restoration of the maxilla (general view and details).

immediately post-operatively, and a pano - ramic tomography every three years thereafter.

The distinct advantage of an orthopantomogram is its panoramic view of all teeth, the osseous limbus alveolaris and important adjacent anatomic structures. By comparison, dental film images as a parallel technique provide information about the progression and stagnancy of the issue degeneration, because

instead, DNA-RNA hybridization tubes were used.

The advantage of these hybridization tubes was that no live material from the probed areas was required for cultivation, thus reducing work in the dental practice. In addition, the results were available much faster than with the classic microbial examination. The disadvantage of these rapid tests is a relatively high price and the fact that the emCASE 1

Fig. 1_Panoramic tomography (emergency service) dating back to 1995-immediately prior to commencement of treatment. Fig. 2-4_Baseline findings in 1995. Fig. 5a & b_Tooth 37 was not conservable in spite of hemisection (August 1995), resulting in a large edentulous space in the third quadrant (November 1995).



- Bone augmentation, if required
- Mucogingival corrections, if required

4. Recall phase (from 05-1995

_After 4 weeks, 6 months, 1 year and then annually: complete survey of clinical evidence, X-ray diagnosis, repeated decontamination with diode laser light of exposed root areas, if required.

_Imaging procedures

As a general rule, the orthopantomogram (pano ramic tomography) and in special cases/as a supplementary measure dental film images as a parallel technique were the applied imaging procedures.

A-scan and B-scan ultrasonography was also used in a few cases of exacerbated inflammations. An orthopantomogram was taken preoperatively and they enable statements about the behavior of the limbus alveo-

Microbial diagnostics

At the time of diagnostic radiology (see above), germ extractions of the affected areas were also performed. This was not done by way of the conventional microbial examination technique (germ extraction— cultivation-pure cultures-microscopic specimen- gas chromatography-sensitivity to antibiotics and color test strips); ployed product only detects special marker germs so that not all microbial organisms in the sulcus can be identified.

The area where a germ extraction was planned had to be carefully dried with a cotton swab. The paper tip was then put in place and, after an exposure time of 10 seconds, was immediately packaged in a sterile container and forwarded to the manufacturer for germ identification. The manufacturer identified the germs and evaluated the socalled marker germ values.

The result was considered negative if less than 0.1 % was identified as a marker germ. The result was considered to be low if 0.1-0.99 % was identified as a marker germ. The result was considered to be medium if 1.0-9.9 % was identified as marker germ and high if more than 10% was identified as marker germ.

Laser light decontamina-

Decontamination was an essential part of the overall therapy: It was achieved with diode laser light of 810 nm wavelength, 1 watt of power and an application time of 20 seconds per tooth and implant under fiber contact

> Fig. 12_The periodontal (vertical bone degeneration) on teeth 15, 14, 24, 25 are so advanced that these teeth can be considered nonconservable.

Fig. 13 & 14_There are essential modifications in comparison with the baseline findings regarding the maxilla. Some teeth have to be extracted. Furthermore, a removable bridge (telescopic bridge) was inserted.









in continuous wave mode. When adhering to these parameters (time limitation and power limitation) it can be guaranteed that the germ spectrum causing the disease can be sufficiently damaged and at the same time that pulpa and/or peri-implant or periodontal tissue structures do not suffer anythermal damage (Bach and Krekeler [1994]).

Three patient cases 1995/2010

Three patients are presented from the original patient group of the "diode laser basic study" (25 from 1995 patients) (Krekeler/Bach, Department of Parodontal Surgery of the Uni-Dental versity Clinic, Freiburg/Breisgau) who showed "typical progression patterns" and whose treatment illustrates the advantage of integrating diode laser light application into a proven therapy regimen for the treatment of marginal parodontopathies.

CASE 3

Fig. 25_X-ray image dating back to 1994 (requested by a previous dentist).

Fig. 26_Panoramic tomography taken in 1995 (commencement of treatment).

Fig. 27 & 28_Initial clinical images dating back to 1995. Fig. 29 Follow-up X-ray image taken

in 2001 (6-year follow-up). Fig. 30_11-year follow-up in 2006. dating back to 1994—prior to commencement of treatment.

Fig. 16_Initial X-ray image taken in 1995.

Figs. 17–19_Baseline findings in 1995.

Fig. 20_Four-year follow-up 1999. Fig. 21_Panoramic tomography taken in 2004; dental implants were inserted to increase the number of abutment teeth.

_1st Case (Figs. 1–14)

The holding therapy case Female patient, born in 1954. Medical history

The patient went to the Sunday emergency service at the Freiburg dental clinic because of pain in tooth 37. A profound parodontopathy was diagnosed there, and the patient came to our department on the following Monday requesting treatment. She had received a complete fixed restoration from her dentist 6 months ago, but without a preprosthetic X-ray diagnosis. Ms. D. is a healthy and very healthconscious physiotherapist.

Clinical baseline findings (1995)

Abutment tooth 17 showed a degree of loosening of 2, as did tooth 26 and tooth 45. Mesial probing resulted in profuse, hard to arrest bleeding. BOP and high probing depths were found in general. The interdental spaces had soft deposits, also under the pontics.

X-ray diagnosis (1995)

The panoramic tomography (orthopantomogram) shows severe horizontal and vertical bone lesions. Teeth 35 and 26 have dish-shaped defects. Trifurcation34 is opened radiologically

Diagnosis

Most severe form of adult marginal parodontitis having portions with a fast-course component.

Course of treatment 1995-2010

Tooth 37 was extracted within the scope of initial pain treatment, as were teeth 26, 17 and 35. Removable immediate prostheses were incorporated because all three pontic reconstructions had to be destroyed during the extraction therapy. The pretreatment phase proved to be unproblematic; the patient was very motivated and eager to learn the oral hygiene techniques as instructed.

cleaning, motivation and instruction steps, a diode laser light application was always performed. Special emphasis was placed on the periodontally severely damaged premolars and the remaining molar 27.

First re-inflammations of the marginal parodontopathy were noticed in 2009; a curettement of teeth 14, 15 and 27 was performed once again. Due to subliminal but latent discomfort, teeth 15, 14 and 27 were removed at the beginning of 2010 and a new concept for treatment of the maxilla was developed.

A removable telescopic prosthesis (cuspids are abutment teeth) was incorporated. The prosthesis on the mandible, which has been in place for 15 years, is still there, and there are no signs of a degeneration of the supportive tissue on the natural and artificial abutment teeth.

Epicrisis

Very remarkable in this patient was the considerable amount of trust she had - in spite of bad experiences in the past in the new laser-assisted therapy concept, which was out of the ordinary at the time. Her compliance was exceptionally good for the entire 15 years. Because of her conscientious oral hygiene and strict adherence to the recall system she remained recur-

_2nd Case (Figs. 15–24)

Success due to (laser-assisted) recall case

Male patient, born in 1938.

Medical history

This patient had been treated since childhood by a dentist who passed away in 1991. For some time the patient had been complaining of too thaches and bleeding of the gums, the latter also occurring spontaneously.

He consulted the successor of his former dentist at the dental practice. However, this dentist did not pay much attention to his descriptions (discomfort) and only remarked once that "there is nothing that can be done!". The patient had obtained the last OPG that had been taken and brought it to the initial examination at our clinic, but he refused (three months later at our clinic) a new

> Fig. 22_The principle of increasing the number of abutment teeth is still being pursued twelve years after commencement of treatment (OPG dating back to 2007). Fig. 23_Current panoramic tomography taken in 2010. Figs. 24_Clinical images taken in 2010: In the frontal view as well as in the lateral views and both jaw views conditions without irritations prevail for the most part.





rence-free for more than a

decade. This still holds true for

the mandible, while the an-

tecedent massive degeneration

of supportive tissue required the

removal of three maxillary teeth.

Thanks to the diode laser as-

sisted periodontal therapy and

the continuous recall, the patient

was able to retain the majority of

her teeth in the maxilla and the

fixed prostheses for a longer pe-

riod of time. It was only recently

that this concept in the maxilla

had to be modified in favor of a re-

movable one; however, this oc-

curred 15 years after a similar

suggestion (removable prosthe-

sis) had been made by her at-

tending dentist at the time.



X-ray diagnosis, stating that he was completely healthy.



Fig.31_The panoramic tomography taken in 2010 does not show any signs of progressive degeneration of the supportive tissue; the image rather shows a "stable osseous

Figs. 32-34_Clinical images 2010 (comprehensive and detail views) showing "reduced" but overall stable conditions. The condition of the mandible also appears normal which is confirmed by the detail view of the frontal mandible.

From June to August 1995 the remaining teeth were treated with open curettement. She had no recurrence for a long time. She received implants in the third quadrant while the remaining maxillary side teeth received fixed prostheses. The edentulous space in the second quadrant remained at the patient's request; in the first quadrant, the principle of a shortened row of teeth was realized (up to 5' to 5th).

tained from the end of 1996 to 2008. The patient conscientiously observed all recall appointments. Aside from the usual

This condition was main-







→ DT page 7

Clinical baseline findings in

Teeth 27 and 37, 38 showed a degree of loosening of I-II. The side teeth showed high probing depths, and a BOP was detected in general. The front mandible was found to be without irritation. The interdental spaces had soft deposits. There were edentulous spaces 16, 25, 26, 27, 45, 46, 35, 36.

X-ray diagnosis (1995)

The panoramic tomography (orthopantomogram) shows an adult dentition with general horizontal bone loss and profound vertical bone lesions on the following teeth: 17,24,27,47 and 48. The patient had received primarily cast restorations. Tooth 24 shows two apical radiopaque structures on the root apex and a discrete periapical translucent zone.

Diagnosis Adult marginal periodontitis.

Course of treatment 1995-2010

The entire pre-treatment phase proved to be without complication due to the patient's initially high compliance. The teeth of the maxilla and the mandible were treated with a mixed open (sidetooth area) and closed (front-tooth area) curettement in the subsequent surgical phase. The surgical part of the periodontal treatment was completed in April of 1995. Since then the patient has been in the recall system, which he took very seriously initially and which helped him to remain recurrence- free for four years after the surgical treatment. From 1999 to 2003 the recall started to become difficult because the patient did not show up in spite of appointments or rescheduled appointments on short notice. At the beginning of 2003 increased probing depth were found on 23, 24 and 27 and three additional teeth exhibited bleeding when probed. Another curettement with laser light decontamination resulted in a decrease of the inflammation; however, 27 could not be saved and had to be extracted, as did tooth 24 (condition after root apex resection), which fractured subgingivally. The resulting freeend situation starting with tooth 23 in the left half of the maxilla and the existing edentulous space in the right half of the maxilla, which had been there for a longer period of time, were treated with three implants that received crowns after a threemonth osseointegration period. We arranged with the patient that he should participate in a quarterly recall and make a new appointment upon completion of the respective recall. He has been recurrence-free since then.

The X-ray images showed a marked tendency for reduction of the osseous supportive tissue on tooth 24. (Note: This tooth was also extracted.) None of the other teeth showed any substantial changes in the course of the osseous limbus alveolaris. The implants also did not show any changes of the periimplant osseous condition from insertion up to the present day.

Epicrisis

Our prognosis after removal of the non-conservable teeth and the systematic increase of abutment teeth is very favorable. The patient's compliance- after variations in the medium observation period— is stable and good. The long recurrence-free interval is also very gratifying.

_3rd Case (Figs. 25–34)

The "completely delightful long-term patient"

Male patient, born in 1952.

Medical history

This patient had been with the same dentist for many years, whom he consulted for checkups on a regular basis. The patient was surprised to find that his teeth 12 and 11 were "loose" and had to be extracted. He was then referred to our clinic. The patient was quite obviously unhappy with the loss of two teeth and the referral ("I feel pushed off"). He is a physical education teacher at a high school and stated that he was completely healthy.

Clinical baseline findings (1995)

Almost all teeth revealed increased probing depths, and probing on the gums in the sidetooth area resulted in bleeding. The smooth surface cleaning was very good; however, deposits were found in the interdental spaces. The dental necks of the maxillary premolars showed wedge-shaped defects. The patient had received primarily cast restorations.

X-ray diagnosis (1995)

In the maxilla, the osseous limbus alveolaris has a considerably reduced level. The alveolar ridge in the area of the tooth gap 12, 11 is severely atrophied. Bone mass in the mandible is also reduced, although not as extensively as in the maxilla. Tooth 45 had received a root canal treatment. The crown edges of the cast restorations do not align perfectly with the contour of the teeth and mostly have an overhanging design.

Diagnosis

Severe adult marginal periodontitis.

Course of treatment 1995 to present

Our most difficult task initially was to appease the patient's dissatisfaction because he felt he "had been taken for a ride." After we had successfully done that, the patient eagerly followed our instructions and followed a frequent and sufficient oral hygiene regime. He grew especially fond of interdental cleaning which had never been mentioned to him before. In May 1995 we started the corrective phase, which was completed in July. We carried out lobe surgery with apical soft tissue fixation in all quadrants. The patient received two implants in regions 12, 11 and, after their osseointegration, two blocked crowns. Due to the severe bone degeneration and the patient's wish to forego augmentation, we arrived far below the cement-enamel junction of the adjacent teeth in one oral implant; however, this did not pose a problem due to the patient's extremely deep-set upper lip. The patient has been in our recall system for 15 years now; he has not missed one recall appointment and has been recurrence-free ever since. A successive prosthetic re-treatment of some single (component) crowns, which had become insufficient, was carried out over the course of several years.

Epicrisis

I feel that—on the "credit side"—we have the patient's excellent cooperation, which has not diminished to this day, and the long recurrence-free period. In this context, one should not forget the extent of the previous periodontitis. These aspects leave a very satisfying impres-

Discussion

The diode laser decontamination study (Krekeler/Bach; University Dental Clinic, Freiburg/Breisgau) that started in 1994 and 1995 was made up of 25 patients, of which seven are still receiving treatment. The extremely long examination period (15 years) naturally limited the number of patients we could examine and treat. Some of the patients, who are no longer in our recall system, have unfortunately passed away in the meantime, while others have moved away or found a different dentist who is located closer to their new residence (usually a care home). Over the years, three "patient types" have emerged-the "holding therapy" type, the "imperative recall type to avoid being the unsuccessful type" and the "successful type". The purpose of this paper is to present these three types by way of individual examples. Diode laser light decontamination proved to be very helpful in all examined patients - Ifeel that, based on the current results, this assessment is justified because the incorporation of diode laser decontamination into the proven treatment regimen for periodontitis resulted in a considerable reduction of the recurrence rate and a considerable improvement of the prognosis of this disease.

An evaluation of the significance of the laser treatment, which has been established as an integral part of a proven therapy regimen in our treatment philosophy, is certainly worthy of discussion. Laser critics will want to argue that a close-meshed and consistent recall, possibly supported by other adjuvant measures, would have yielded similarly positive results. This may indisputably be the case; in fact, I am sure that this assessment is true! However, if the key to treatment success is then rather the consistency and frequency of treatment, I consider laser-assisted treatment to be one of many options in the extensive field of periodontal therapy. Laser-assisted periodontal therapy thus makes no pretence of being a unique feature, but rather an adjuvant therapy with the claim to be efficient, gentle and ultimately successful._

AD



EXPERT FACULTY

DIDIER DIETSCHI, MAURO FRADEANI, DANIEL EDELHOFF, STAVROS PELEKANOS, PANOS BAZOS, ED MCLAREN, SASCHA JOVANOVIC, PASCAL & MICHEL MAGNE, BRIAN LESAGE & OTHERS ...



Register & More Information at:

Contact in Athens: O### Tel: +30 210 213 2084, +30 210 222 2637 E-mail: lito@omnipress.gr, omnip

Web:www.omnicongresses.gr Contact in the US:

Nena Puga Tel.: +1 310 696 9025 gIDE GLOBAL INSTITUTE E-mail: nena@gidedental.com website: www.gidedental.com

Media Partner: dti]

(W&H)

SPONSORS

STHETIC CURRICULUM

Dr. Ed McLaren & Dr. Sascha Jovanovic

Dr. Didier Dietschi, Dr. Daniel Edelholff, Dr. Mauro Fradeani

SESSION I - FEBRUARY 18 - 21, 2011 IN ATHENS, GREECE

DIRECT RESTORATIONS AND ADHESIVES with DR. DIDIER DIETSCHI and others

SESSION II - JUNE 24 - 27, 2011 IN ATHENS, GREECE

INDIRECT RESTORATIONS, FULL CERAMIC **CROWNS AND VENEERS**

SESSION III - OCTOBER 2 - 5, 2011

with DR. DANIEL EDELHOFF and others

IN ATHENS, GREECE

ALL CERAMIC RESTORATIONS AND **FIXED PROSTHODONTICS** with DR. MAURO FREDEANI & DR. STAVROS PELEKANOS

SESSION IV - NOVEMBER 28 - DECEMBER 2, 2011 N LOS ANGELES, CALIFORNIA

GRADUATING WEEK AT UCLA with DRS. ED MCLAREN, SASCHA JOVANOVIC, BRIAN LESAGE TODD SCHOENBAUM, PASCAL & MICHEL MAGNE

PROGRAM FEATURES

- CLASSROOM EDUCATION
- LIVE PATIENT TREATMENTS
- CASE PRESENTATIONS
- HANDS-ON
- 90 HOURS SELF-STUDY
- 217 HOURS CE
- 2 CERTIFICATES from gIDE and UCLA





DTSC - COURSES, COMMUNITY, TECHNOLOGY, ON-DEMAND

The DT Study Club makes all of this possible from the comfort of your own computer and without travel expenses. In other words, welcome to the community!

The purpose of this study club is to provide practitioners like yourself an opportunity to learn and network with like-minded colleagues in a friendly, non-threatening environment. We encourage you to take advantage of Dental Tribune's global outreach to access a variety of fresh perspectives and cultures, enhancing your educational mix.





focused on helping today's practitioners to stay up to date. Networking possibilities that go beyond borders to create a truly Global Dental Village



VIDEO REVIEWS OF PRODUCTS

Our opinion leaders unveil new products, services, and give you their first impressions of the industry's hottest topics.



PEER REVIEWED **CASE STUDIES**

Upload, comment, participate. We encourage you to share your cases for review with like-minded practitioners.





REGISTER FOR FREE ON WWW.DTSTUDYCLUB.COM

CONTESTS WITH CHANCES TO WIN FREE TUITION FOR ADA/CERP C.E. ACCREDITED WEBINARS

SPONSORSHIP AND SPEAKING INQUIRIES: JULIA WEHKAMP, J.WEHKAMP@DTSTUDYCLUB.COM, (416) 907-9836.



















ADA C·E·R·P® Continuing Education Recognition Program



DENTAL HYGIENE



