

DENTAL TRIBUNE

—The World's Dental Newspaper • Middle East & Africa Edition—

PUBLISHED IN DUBAI


APRIL 2009

No. 2 Vol. 7



News
April 26: International
Seeds Day

Page 3



Media CME
TM Disorders: Treat-
ment & Management
Considerations

Page 6



Trends & Applications
Analysis of Mercury in
Wastewater

Page 11

Gum disease linked to gestational diabetes

A new study by NYU dental researchers has uncovered evidence that pregnant women with periodontal (gum) disease face an increased risk of developing gestational diabetes even if they don't smoke or drink, a finding that underscores how important it is for all expectant mothers - even those without other risk factors - to maintain good oral health.

The study, led by Dr. Ananda P. Dasanayake, Professor of Epidemiology & Health Promotion at New York University College of Dentistry in collaboration with the Faculty of Dental Sciences at the University of Peradeniya, Sri Lanka, eliminated smoking and alcohol use among a group of 190 pregnant women in the South Asian island nation of Sri Lanka, where a combination of cultural taboos and poverty deter the majority of women from smoking and drinking. The findings support an earlier study led by Dr. Dasanayake that found evidence that pregnant women with periodontal disease are more likely to develop gestational diabetes

than pregnant women with healthy gums.

That study, which followed 256 women at New York's Bellevue Hospital Center through their first six months of pregnancy, showed that 22 of the women developed gestational diabetes. Those women had significantly higher levels of periodontal bacteria and inflammation than the other women in the study. The findings were published in the April 2008 issue of the Journal of Dental Research.

More than one-third of the women in the new study, which was conducted over the course of one year, reported having bleeding gums when they brushed their teeth. The women were given a dental examination and a glucose challenge test, which is used specifically to screen for gestational diabetes. According to Dr. Dasanayake, those women found to have the greatest amount of bleeding in their gums also had the highest levels of glucose in their blood. Dr. Dasanayake, who presented the findings today at the annual meeting of the International As-

sociation for Dental Research in Miami, said that he expected the final data to show that between 20 and 30 of the women had developed gestational di-

abetes.


Gestational diabetes is characterized by an inability to transport glucose -- the main source of fuel for the body -- to the cells during pregnancy. The condition usually disappears when the pregnancy ends, but women who have had gestational diabetes are at a greater risk of developing the most common form of diabetes, known as Type 2 diabetes, later in life. Asians, Hispanics, and Native Americans are at the highest risk for developing gestational diabetes. All of the women in the Sri Lanka study were of Asian origin, while 80 percent of the New York study subjects were Hispanic.

"In addition to its potential role in preterm delivery, evidence that gum disease may also contribute to gestational diabetes suggests that women should see a dentist if they plan to get pregnant, and after becoming pregnant," Dr. Dasanayake said. "Treating gum disease during pregnancy has been shown to be safe and effective in improving women's oral health and minimizing potential risks." [D]

Our ears may have built-in passwords

People may no longer need to remember their password to access bank accounts, they can do that by picking up a phone and letting their ear do the talking, Researchers in England are developing a new biometric technique that elicits the sound ear hairs create when ruffled by noise. If each person's "ear sound" is unique, and stays the same over time, it could become a high-tech password to access accounts and cell-phones.

That's because "hearing is an active process - the ear actually puts energy into the incoming sound waves to replace energy lost as sound is absorbed by the ear's structure", says Stephen Beeby, an engineer at the University of Southampton, UK, who is leading the research. "This



process helps us hear things we otherwise would not.

Predicted in the 1940s but only confirmed with ultralow-noise microphones in the 1970s, ear-generated sound is evoked with a series of clicking noises. Anecdotal, experts say they can differentiate one ear from another, but it "has to be able to reliably recognize people over long time periods," one scientist said. "For example, a fingerprint taken from a 20-year-old is still valid when they are 60." [D]

PROMEDICA

Highest quality made in Germany

Nano-Ceram-Technology

- ▶ nano-reinforced ceramic particles
- ▶ special resin matrix

outstanding biocompatibility

excellent aesthetics:

- ▶ exceptional aesthetics and polish
- ▶ high colour stability
- ▶ perfect translucency and colour adaptation

excellent physical properties:

- ▶ durable, long-lasting restorations
- ▶ perfect permanent marginal tightness
- ▶ extremely low polymerisation shrinkage
- ▶ optimal abrasion resistance

PROMEDICA Dental Material GmbH

Tel. +49 43 21 / 5 41 73 · Fax +49 43 21 / 5 19 08

Internet: <http://www.promedica.de> · eMail: info@promedica.de



flowable version



Light-curing nano-ceram composite

- universal for all cavity classes
- comfortable handling, easy modelling

- exact match of translucency and shade to the packable Composan bio-esthetic

PROMEDICA

AD

PARIS: People who drink their tea piping hot run a higher risk of throat cancer than counterparts who prefer a cooler cuppa, according to an investigation published on Friday by the British Medical Journal.

factor, possibly because of damage to throat tissue. Interested in finding out more, Iranian researchers went to Golestan province, which has one of the highest rates of oesophageal cancer in the world.

A team led by Reza Malekzadeh of the Digestive Disease Research Centre at Tehran University of Medical Sciences looked at 300 people who had been diagnosed with a throat tumour and a matched group of 571 healthy people who lived in the same area.

compared with those who drank warm or lukewarm tea, whose temperature was 65°C or less. Drinking very hot tea (at least 70°C) was associated with an eightfold increased risk compared with warm or lukewarm tea.

actly how this happens remains unclear.

Though some might see sports drinks as a healthier alternative to soda, a new study shows that the citric acid they contain can damage teeth.

The finding comes from a study involving teeth from cows. New York University College of Dentistry researchers cut the teeth in half and placed them in top-selling sports drinks. After soaking for up to 90 minutes, which the researchers said simulated sipping on the drinks throughout the day, the enamel coating of the teeth was partially eaten away. This allowed the drinks to leak into the bonelike material underneath the enamel, causing the teeth to soften and weaken.

The condition, called erosive tooth wear, can result in severe tooth damage and tooth loss, if not treated.

"This is the first time that the citric acid in sports drinks has been linked to erosive tooth

wear," study leader Dr. Mark Wolff, chairman of cariology and comprehensive care at the NYU College of Dentistry, said in a news release issued by the school.

The findings were to be presented Friday at the International Association for Dental Research general sessions in Miami Beach, Fla.

Perhaps surprisingly, brushing immediately after having a sports drink might actually cause more damage, Wolff said, as the softened tooth enamel is vulnerable to the abrasiveness of toothpaste.

"To prevent tooth erosion, consume sports drinks in moderation and wait at least 30 minutes before brushing your teeth, to allow softened enamel to re-harden," he said. "If you frequently consume sports drinks, ask your dentist if you should use an acid-neutralizing, re-mineralizing toothpaste to help re-harden soft enamel." [DT](#)

Scientists have used tobacco to produce medicine for the treatment of a number of diseases, including diabetes.

Research published in the journal BMC Biotechnology details how genetically modified tobacco plants were used to produce medicines for several autoimmune and inflammatory diseases.

Experts set out to create transgenic tobacco plants that would produce biologically-active interleukin-10 (IL-10), a potent anti-inflammatory cytokine.

They tried two different versions of IL-10 (one from a virus, one from the mouse) and generated plants in which this protein was targeted to three different compartments within the cell, to see which would work most effectively.

The researchers found tobacco plants were able to process both forms of IL-10 correctly, producing the active cytokine at high enough levels that it might be possible to use tobacco leaves without lengthy extraction and purification processes.

The authors claim they are keen to use the plants to see whether repeated small doses could help prevent type 1 diabetes mellitus (T1DM), in combination with other auto-antigens associated with the disease.

Commenting on the research, Professor Mario Pezzotti from the University of Verona, said: "Transgenic plants are attractive systems for the production of therapeutic proteins because they offer the possibility of large scale production at low cost, and they have low maintenance requirements.

"The fact that they can be eaten, which delivers the drug where it is needed, thus avoiding lengthy purification procedures, is another plus compared with traditional drug synthesis." **DT**

Prof. Abdullah Al-Shammary, Restorative Dentistry, KSA
Prof. Hussain F. Al Huwaizi, Endodontics, Iraq
Prof. Samar Burgan, Oral Medicine, Jordan
Dr. Abdel Salam Al Askary, Implantology, Egypt

Dr. Talal Al-Harbi, Orthodontist, Qatar
Dr. Mohammed H. Al Jishi, Bahrain
Dr. Lara Bakaeen, Prosthodontist, Jordan
Dr. Aisha Sultan, Periodontist, UAE
Dr. Kamal Balaghi Mobin Aesthetics, Iran

The World's Dental Newspaper • Middle East & Africa Edition

Published by Venus Advertising

in licence of Dental Tribune International GmbH

© 2008, Dental Tribune International GmbH. All rights reserved.

Dental Tribune makes every effort to report clinical information and manufacturer's product news accurately, but cannot assume responsibility for the validity of product claims, or for typographical errors. The publishers also do not assume responsibility for product names or claims, or statements made by advertisers. Opinions expressed by authors are their own and may not reflect those of Dental Tribune International.

President/CEO

Yasir Allawi
y.allawi@dental-tribune.ae

Director mCME:

Dr. D. Mollova
info@cappmea.com

Marketing manager

Sawsan Alhalwachi
info@dental-tribune.ae

Production manager

Hussain Alvi
hussain.alvi@dental-tribune.ae

P.O. Box 4473, Dubai, United Arab Emirates,
Tel. +9714-2282928, Fax: +9714-2241024
E-mail: info@dental-tribune.ae



April 26: International Seeds Day

Order 81: "Iraqi farmers shall be prohibited from re-using seeds of protected varieties,"

Organizations, activists and people from various professional and linguistic backgrounds will observe April 26 as International Seeds Day (ISD) advocating for patent-free seeds, organic food and farmers' rights. ISD will be an educational day for the public to learn about genetically modified food and its health hazardous effects and the agribusiness of major US and European companies and their monopoly over the agriculture in Africa and Asia with emphasis on India, Iraq and Afghanistan. It will be a day of solidarity with farmers in countries devastated by war (Afghanistan, Iraq & others) and of resistance: <http://www.INEAS.org/events.htm>

Historically, the Iraqi constitution prohibited ownership of biological resources. Farmers in Iraq have operated in a mostly free-to-little-regulated, informal seed supply system. Farm-saved seeds and the free exchange of planting materials among farmers have long been the basis of agricultural practice in Iraq. Yet all of this has become history.

On April 26, 2004, Paul Bremer, the administrator of the Coalition Provisional Authority (CPA), issued and signed Order 81, which prohibits farmers from reusing seeds harvested from new varieties registered under the law. When ownership of a crop is claimed, seed saving will be banned and farmers will have to be pay royalties to the registered, so-called seed owner.

Order 81 was signed on April 26, 2004 by Paul Bremer, the administrator of the Coalition Provisional Authority (CPA) in Iraq to control Iraq's agriculture. The Order was a declaration of war against farmers. Article 14 of this law states "Farmers shall be prohibited from re-using seeds of protected varieties," Order 81 mends Iraq's original law No. 65 on patents, created in 1970.

A "Greedy law, unjust law is meant to be disobeyed" <http://www.youtube.com/watch?v=I-B1yU278z>

The most significant part of Order 81 is the subject of 'Plant Variety Protection' (PVP), which ensures not the protection of biodiversity, but rather the protection of the commercial interests of USA and European major seed corporations. In order to qualify for PVP, seeds have to be 'new, distinct, uniform and stable'. Therefore, the sort of seeds being encouraged to grow by corporations such as World Wide Wheat Company (WWWC), Monsanto and others will be those registered under PVP.

The law awarded US Corporations complete control over farmers' seed for 20 years. Iraqi farmers had to sign an agreement to pay a "technology fee" plus an annual license fee. Plant Variety Protection (PVP) made seed reusing and saving illegal as well as "similar" seed plantings punishable by severe fines and imprisonment. Agribusiness wants the same rights everywhere, including in the USA. This

will jeopardize the future of organic and independent farming.

According to Latha Jishnu, senior editor of Business Standard "Five years after Order 81 was passed, farm activists across the world have got together to mark April 26 as International Seeds Day to help Iraqi farmers to break the

vice-like grip of the global seed companies. The campaign is coordinated by the Institute of Near Eastern & African Studies (INEAS), based in Cambridge, Massachusetts, and has got the backing of some organisations in India. I met Wafaa Al-Natheema of INEAS, when she was in India earlier in the year to drum up support for the

campaign, and she says the world needs to respond to this threat to agriculture. Iraqi farmers, like the rest of the nation, are unaware of this law and how it could turn their world upside down. "They need our help to learn how to retain their seeds under these circumstances and how to lobby against this unjust law." <http://www.business-standard.com/india/news/latha-jishnu-order-81-the-plunder-farming/353518/>

For more information:

The Institute of Near Eastern & African Studies (INEAS)
1 (617) 86-INEAS (864-6327)
INEAS@aol.com INEAS_1994@yahoo.com Full text of Order 81: http://www.trade.gov/static/iraq_memo81.pdf
People's rights to water and food: <http://www.navdanya.org/organic/index.htm>
The agenda in Afghanistan & Iraq: <http://www.grain.org/briefings/?id=217>

AD



If it isn't clean, it can't be sterilised!
Hydrim and Statim, your perfect partners
in the sterilisation process

Now you can eliminate the risk of hand washing instruments

Authorities now recommend that instruments cannot effectively be sterilised unless they have first been cleaned in a mechanical washer. Across Europe guidelines are under review to reflect this advice.

That's why effective sterilisation begins with the Hydrim® C51wd or M2 instrument washer. After a pre-wash to remove virtually all proteins, two independent, high pressure sprays clean the instrument load from above and below, removing virtually all organic debris. Independent tests show an efficacy of 99.9 – 100 per cent. The Hydrim washers perfectly prepare instruments for sterilisation in as little as 19 minutes, and eliminate the risk of washing instruments by hand.

Washed and
sterilised in
27 minutes

Following the Hydrim wash cycle, the instrument basket can immediately be transferred into an S class Statim® autoclave cassette for the fastest sterilisation cycle available. Most instruments, including handpieces can now be sterilised between patients in a Statim 2000 S in only 8 minutes. The fast Statim S cycle process fully complies with the latest European standard EN13060.

The compact Hydrim C51wd and Statim units are designed for benchtop use and are the perfect partnership for the central sterilisation area even in the smallest practice. The floor standing Hydrim M2 can process about 100 instrument sets in a 10 hour day and is ideal for the larger, busy practice.

For further information about SciCan products please contact:

Sub Sahara and Africa – Eric Bernard on +27 11 486 1622, +27 82 788 7817, or email ebarnard@global.co.za

Middle East – Ashraf Suleiman on +20 12 326 7870 or email asuleiman@scican.com or visit our website at www.scican.com

SciCan
A HIGHER STANDARD

American Medical Association may investigate journal editors

CHICAGO (AP) — The American Medical Association is seeking an investigation of claims that editors of its leading medical journal threatened a whistleblower who pointed out a researcher's conflict of interest.

Editors of the Journal of the American Medical Association deny threatening a professor who raised concerns about a study author's undisclosed financial link to a drug company

when JAMA published the study last year. JAMA, like most leading medical journals, has a policy of noting scientists' industry connections.

According to the Wall Street Journal, JAMA editors threatened to ban the professor from their journal and ruin his medical school's reputation if he didn't stop talking to reporters.

The editors deny that. But the flap prompted them to spell out

what amounts to a gag order on anyone who alerts the medical journal about suspicions that a researcher has undisclosed industry ties. The journal editors argue that any suspicions should be kept secret until JAMA can complete its own probe. That is an existing policy, JAMA's editor-in-chief, Dr. Catherine DeAngelis, told The Associated Press on Monday. AMA journals are independent and the medical associ-

ation doesn't interfere with what they publish. But AMA said Friday it has asked an independent oversight committee to investigate how JAMA editors handled the issue.

"As owner and publisher of JAMA, we take these concerns very seriously," AMA board chairman Dr. Joseph Heyman said in a written statement.

The issue involves a study published in JAMA last May that

said the drug Lexapro prevents depression in stroke patients. A Tennessee university professor who reads JAMA told the editors in October that he had learned that a study author had served as a speaker for Lexapro's maker. Though other industry ties were noted in the journal, that one was not. JAMA editors vowed to investigate. The professor, Jonathan Leo of Lincoln Memorial University, also discussed his concerns in a March 5 letter posted on a different medical journal's website.

On March 11, JAMA editors published a correction revealing the undisclosed ties to Lexapro's maker. JAMA's editors acknowledged in a March 20 editorial being upset about Leo airing his concerns. They argue that publicizing unconfirmed allegations about study authors could unfairly damage reputations and interfere with JAMA's own investigations. [D1](#)

Confusion Over Osteoporosis Medication and the Affect on Oral Health

It has been estimated that over 10 million Americans have Osteoporosis; a condition that causes the thinning of the bone along with severe loss of bone density over time. Medications, specifically bisphosphonate are used to treat this often debilitating disease. Although these drugs effectively slow bone loss and possibly increase bone density reducing the risk of fracture, they have recently come under-fire as reports have suggested the use of medications containing bisphosphonates may cause individuals to develop "bisphosphonate-associated" osteonecrosis of the jaw. Considered to be a rare condition osteonecrosis of the jaw results in severe damage due to the temporary or permanent loss of blood supply to the jaw bone possibly causing pain, numbness, exposed bone, tooth loss, and infection.

The American Dental Association and the National Osteoporosis Foundation have teamed-up to provide patients with a brochure titled "Osteoporosis Medications and Your Dental Health," that will be available from dental offices this month. The goal is to separate fact from fiction by providing the brochure as a resource to patients that have become alarmed by the suggestion that their medication may be causing osteonecrosis of the jaw. In a press release Matthew Messina, D.D.S., ADA Consumer Advisor states that "Patients who take bisphosphonates for osteoporosis are encouraged to talk to their dentist so that their dentist can show them good oral hygiene practices as well as monitor their oral health," adding stress to the point that "Patients should not stop taking their osteoporosis medications without speaking with their physicians." [D1](#)

FDI Annual World Dental Congress 2-5 September 2009 Singapore



congress@fdiworldental.org
www.fdiworldental.org

fdi
SINGAPORE 2009

DTME talks to Mr. Andrew Bennett, Business Developer for 3M Unitek , to find out his views about the Middle East Market and current economic situation

3M Unitek Orthodontist Symposium Transcription

What are the objectives of this Symposium?

The aim is to challenge the traditional core process and to put out some new ideas as to where we think the industry will be going over the next 5, 10 or 15 years. This is not something we would do every single year because the developments achieved are so far advanced that the industry doesn't change enough to warrant 2 days here in beautiful Dubai every year.

The fact that all these orthodontics are still here and didn't cancel their registration proves that they are all still interested in these new cutting-edge concepts from 3M Unitek.

It is quite unusual to be able to get this many orthodontics from across the world together all at the same time. Usually at dental conferences you would have a big group of general dentists so it is great to see so many specialist orthodontists, especially with the current economic climate.

How is the current economic climate impacting 3M Unitek right now?

Of course, this is an elective treatment so – ultimately its not like heart surgery or something ... you have straight teeth or you don't. I think what we are seeing to a small extent, and I literally just relocated from the US, where the economic environment is very different, we saw doctors tending to purchase products when they needed them rather than build up a large inventory. From our perspective we have seen no slow down at all in this region.

We acquired the Incognito brand last year which is a very high end treatment in the Lingual side of the Market. They continue to be a huge success in this region and certainly in mainland Europe this is probably the most expensive treatment you can have. It continues to be a great success and a great acquisition for us. Even as we operate on the higher end of the market everything seems to be ok – the proof of having 500 doctors being prepared to travel from 40 countries here kind of says business is good - business is very buoyant.

As I mentioned earlier, it is an elective treatment so we always know there is always a risk with any treatment which is elective – it's not bandages and it's not tape, but its orthodontics.

You're new to the region, Middle East and Europe –How are you looking forward in this market?

I have been with 3M and Unitek for 15 years and was in the US before I came here. The 3M Unitek range operates from the very top of the pyramid to the very bottom in terms of price perspectives and quality perceptive - our intention is to be able to cover the broadest possible range of doctors as we can and we have done that with acquisitions recently.

Incognito is proof of that. I think you will that as we go forward and expand our range will broaden considerably.

3M Unitek is a business which likes to operate directly with its customers – in 3M's world this is quite unusual. 3M tends to be a distributor to channel based busi-

nesses, 3M Unitek is one of 2 businesses in 3M that is almost entirely direct. It enables us, I think, to service the customers much better. I think we have 50,000 customers, 14,000 SKU's – which is huge for 3M – we will keep driving that model forward, because

it just gives us a better relationship with our customers.

Ultimately our plan has been for many years to build and sustain the best relationships in the

→ DT page 14



Mr. Andrew Bennett
BDM for EMEA – 3M UNITEK

AD



Enhance your patients' confidence with the range of COREGA denture Fixatives and Cleansers.

Benefits of COREGA denture Fixatives:

- Forms a protective cushion to keep irritating food particles out
- Increases the bite force up to 5 times even for well fitting dentures
- Provides increased denture retention and stability

Benefits of COREGA denture Cleansers:

- Eliminates 99.9% of the odour causing bacteria
- Cleans areas that brushing can miss
- Removes plaque, stains and food debris
- Freshens breath



SPEAK, EAT AND SMILE WITH CONFIDENCE



TM Disorders: (Part 3 of 3)ⁱ

Treatment & Management

Considerations

Ulises A. Guzman & Henry A. Gremillion, USA

The article has been accredited by Health Authority - Abu Dhabi as having educational content and is acceptable for up to 2 (Category 1) credit hours. Credit may be claimed for one year from the date of subscription.

Temporomandibular disorder (TMD) represents a multiplicity of conditions expressed in the masticatory system affecting the temporomandibular joints, masticatory muscles, and/or the associated structures. Many of these conditions share common signs and symptoms, yet require differing treatment/management approaches. Therefore, it is important to identify the specific subcategory of TMD in order to develop a case-specific plan of care.

In addition, etiologic variables and factors associated with perpetuation or recurrence of TMD must be appreciated and determined for each patient. A complete evaluation of each case from historical, clinical presentation and physical/psychological perspectives must be accomplished. Treatment outcomes can be enhanced by the identification of and management strategies that address all the components involved.

The development of a diagnosis-specific plan with a prioritized problem list is necessary to enhance our treatment prognosis. The primary goals of treatment of TMD are to: reduce or eliminate pain; restore a more normal function; allow return to the activities of daily living; reduce long-term health care needs for the problem.

A multi-disciplinary model that includes patient education and self care, cognitive behavioral intervention, pharmacotherapy, physical therapy and orthopedic appliance therapy (interocclusal splints) is favored for the management of the vast majority of TMD patients. It is important to understand that the natural course of TMD does not reflect a progressive disease process, but rather TMD appears to be a complex disorder that is affected by a multitude of interacting factors serving to maintain the disorder or result in recurrence.¹⁻⁵

Most TMD patients will obtain significant improvement of signs and symptoms with a conservative model (non-surgical modalities). Many studies have supported that most TMD patients have minimal or no symptoms after treatment with conservative therapy.⁶⁻⁸ Studies related to intracapsular disorders have demonstrated that in patients with disc displacement (with or without reduction), the natural progression of the disease can

allow for changes that are favorable for a significant number of patients in terms of function and symptoms.⁹⁻¹¹

Involving the patient in the physical and behavioral management of his/her condition is essential in the treatment outcome. As clinicians in the development of an individualized plan of management, we must determine if intervention is necessary, if the condition is acute or chronic and what would be the prognosis of the condition with and without treatment. If intervention is in the patient's best interest, then we must determine to what degree should we intervene (reversible versus irreversible treatment) and decide between a monodisciplinary versus a multidisciplinary approach.

Patient education and self-care is based on the patients' knowledge of their pain concern. The aim of a self-care program is to prevent further injury to the musculoskeletal system and to allow for a period of healing to take place. The success of self-care depends on patient motivation, cooperation and compliance.¹ The most important aspect of self-care is ongoing encouragement and reinforcement by the clinician.

Self-directed care typically includes: limitation of mandibular function, habit awareness and modification, a home exercise program and stress management. Promoting rest for the injured tissues promotes healing. Voluntary limitation of mandibular function, maintaining a soft diet, avoidance of foods that require a great deal of chewing, opening wide, yawning or other activities that promote excessive mandibular function should be avoided.

Clenching, bruxism, maladaptive tongue position habits and other habitual behaviors must be identified. Correction or behavior modification may require clinical assistance.^{12,15} An individualized home exercise program with a detailed description of the program will not only enhance the doctor-patient relationship, but will also assure the patient's compliance, thus making treatment more effective and resulting in a faster rehabilitation.

A program of moist heat and/or ice to the affected areas, massage of the affected muscles, and controlled mandibular movement can enhance joint lubrication and nutrition by encouraging the production of physiological quality and quantity of synovial fluid and minimizing the accumulation of metabolic

by-products and pain mediating substances.

Identification of the source(s) of stress and the importance of the patient understanding the association and adverse influence of stress and the course of TMD are also vital. Clinical and health psychologist participation in your multidisciplinary approach may be required to enhance your treatment outcome.

Pharmacotherapy

Rational utilization of pharmacological agents can be a valuable adjunct in the treatment of TMD. Drugs must be considered on a case-specific basis. A clinician must remember that the treatment of TMD cannot rely on a single drug for all cases. Understanding the variety of drugs utilized in the treatment of musculoskeletal conditions, their potential drug interactions and their side effects can result in a useful tool in our armamentarium.

The most effective pharmacological agents for the management of TMD include analgesics, non-steroidal anti-inflammatory drugs (NSAIDs), corticosteroids, anxiolytics, muscle relaxants, and antidepressant at very low dosages.^{14,15}

Non-steroidal Anti-inflammatory Drugs

This category is effective for the management of mild to moderate pain and inflammatory conditions, particularly those of muscle origin. Relief of symptoms is typically achieved prior to the anti-inflammatory effect. In order to obtain anti-inflammatory effects, these medications should be taken for a minimum of two weeks following the recommended schedule. NSAIDs differ in formulation, efficacy and toxicity. It is suggested that if one NSAIDs fails, another agent should be considered. Common side effects to be considered include gastric distress, inhibition of platelet aggregation, tinnitus/dizziness, and renal/liver toxicity. A list of the most commonly NSAIDs utilized is found in Table 1.

Steroids

Corticosteroids are typically indicated in cases of non-infectious inflammation when NSAIDs have proven to be ineffective. Systemic corticosteroids are not commonly prescribed in the treatment of TMD due to their side effects. They could be considered when in association with the polyarthrides. Intra-articular temporomandibular injection of corticosteroids has been recommended on a selective and limited basis in cases of severe joint pain or in cases of flare ups where conservative therapy has failed.^{16,17} We must recognize that multiple intracapsular steroid in-

Category	Generic	Brand	Dose (mg)
Salicylates	ASA	Bayer	q4h (300)
	Salsalate	Disalcid	bid, tid (500)
	Diffunisal	Dolobid	bid, tid (500)
Propionic Acid	Ibuprofen	Motrin	tid, qid (600-800)
	Naproxen sodium	Naprosyn	qid 375, bid (500)
Acetic Acid	Indomethacin	Indocin	tid (25-50)
COX 2 Inhibitors	celecoxib	Celebrex	Qd, bid (12.5-25)

Table 1: Non-steroidal anti-inflammatory drugs

(Methylprednisolone) Medrol Dosepack	4 mg tablets
(Betamethasone) Celestone	6 mg/cc
(Dexamethasone) Decadron	4 mg/cc

Table 2: Steroids

Generic	Brand	Dose (mg)
Diazepam	Valium	2-5 mg tid
Clonazepam	Klonopin	.05-1 mg tid
Lorazepam	Ativan	0.5-1mg tid
Temazepam	Restoril	15-30mg qhs

Table 3: Antianxiety agents

Generic	Brand	Dose (mg)
Carisoprodol	Soma	350 mg tid
Methocarbamol	Robaxin	750 mg tid
Cyclobenzaprine	Flexeril	10 mg tid
Diazepam	Valium	2-5 mg tid

Table 4: Muscle Relaxants

Generic	Brand	Dose (mg)
Amitriptyline	Elavil	10-75
Desipramine	Norpramin	10-50
Nortriptyline	Pamelor	10-75
Doxepin	Sinequan	10-75

Table 5: Antidepressant agents

jections may have detrimental effects.

These medications are also effective in the treatment of inflammatory conditions such as tendonitis or tendomyositis where, due to the decreased blood flow to the areas, oral medications will provide less than desirable results. Side effects include decreased resistance to infection, fluid retention weight gain, painless myopathy, suppression of the hypothalamic-pituitary-adrenal (HPA) axis, osteoporosis and mood alteration with only short term use. Steroidal medications commonly used are listed in Table 2.

Anxiolytics

Anti-anxiety medication may be utilized as supportive therapy in cases where high levels of emotional stress are associated with TMD. Diazepam can be prescribed for acute exacerbation of masticatory muscle pain,⁵ sleep disturbances and moving disorders such as bruxism.^{14,18} Due to the significant potential for dependency and/or addiction, these

medications should only be used on a short term basis. No more than a 10-day consecutive period when utilized multiple times a day, and no more than three weeks when utilized at bedtime.

Klonopin has shown to have a significant effect in patients with myofascial pain.¹⁹ Side effects include drowsiness and nausea. Benzodiazepines are contraindicated in patients with narrow-angle glaucoma, and can increase CNS depression. A list of anxiolytic agents typically utilized in TMD, sleep disturbances to include insomnia, and moving disorders such as bruxism are included in Table 3.

Muscle Relaxants

Centrally acting muscle relaxants are frequently used in the treatment of temporomandibular disorders.²⁰ It is still uncertain the mechanism in which the benefit from this medications is obtained due to the clinical efficacy at low doses. Either due to their selective effect on relieving muscle spasm

ⁱ Part 1 appeared in *Dental Tribune Asia Pacific*, No. 9 Vol. 5, September 2007; Part 2 appeared in *Dental Tribune Asia Pacific*, No. 10 Vol. 5, October 2007.

or due to their action as a sedative, they play an important role in the treatment of TMD. Primary indications are for muscle spasm, acute muscle pain and to help prevent the increased muscle activity associated with TMD.

Flexeril (cyclobenzaprine hydrochloride), which is similar chemically to tricyclic antidepressants, is the drug of choice for generalized chronic muscle pain. Flexeril has been shown to provide significant relief of muscle pain, and enhance the quality and quantity of sleep. Its combination with an NSAID can be a very effective tool in the treatment of acute TMD. Diazepam, a benzodiazepine is also used as a muscle relaxant. A list of commonly used muscle relaxants is shown in Table 4.

Antidepressants

These medications are helpful with chronic diffuse pain due to myofascial pain, especially when it has been recognized that sleep disturbance is a contributing factor. The analgesic properties of the tricyclic antidepressants are independent of the antidepressant effect. They have shown pain modification properties at therapeutic dosages much lower than those prescribed for antidepressant effect.

The therapeutic effect of the drugs is thought to be related to their ability to increase the availability of the neurotransmitters serotonin and norepinephrine at the synaptic junction in the central nervous system. Studies have demonstrated their use also in the treatment of sleep related bruxism, tension type headache, migraine headache prophylaxis, fibromyalgia and various neuropathic conditions.^{21,22}

Side effects are mainly related to the anticholinergic activity that induces xerostomia, constipation, fluid retention and weight gain. Patients occasionally complain of sedation upon awakening. Contra-indications include cardiac arrhythmias, seizure disorders and patients suffering from panic attacks. Dosages should begin at the lowest level (10 mg) at bedtime and be increased each week only if needed and tolerated by the patient. Table 5 shows a list of some of the most commonly utilized drugs in this class.

Opioids

Typical indications for opioids in the TMD population include exacerbation of pain, postoperatively and in cases of overt trauma. These medications are best indicated for moderate to severe pain over a short period of time. Most common side effects are nausea, respiratory depression and physical dependence. Opioids may be considered in cases of pain refractory for appropriately integrated multidisciplinary care when properly monitored.

Local Anesthetics

Local anesthetics can be useful in the TMD population as a diagnostic tool and also in selective cases as a therapeutic modality.

Indications are as a diagnostic block and in the management of myofascial trigger points. Injections into skeletal muscle with local anesthetics that contain a vasoconstrictor can increase the toxicity of the solution.

Typically, lidocaine or carbocaine without a vasoconstrictor is recommended, especially when injected into muscle (to minimize myotoxic effects). Diagnostic anesthesia may be as simple as the usage of a topical agent, somatic blocks (infiltration,

field blocks and division blocks), trigger points injections, temporomandibular joint injections and/or a sympathetic neural blockade.

Physical Therapy/Physical Medicine

The goal is to relieve musculoskeletal pain, restore normal function, reduce inflammation, coordinate and strengthen muscle activity and promote repair and regeneration of tissues. Rehabilitation of the compromised masticatory system may require

various physical techniques.⁵ Close cooperation with a physical therapist/physical medicine practitioner who is well trained in the management of musculoskeletal disorders of the head and neck is essential.²²⁻²⁵

Massage

Massage over the painful areas is thought to produce an alteration in the sensory input that exerts an inhibitory influence on pain. It is used to reduce edema and to increase blood flow to the area.

Joint Mobilization

The goal is to passively restore joint motion and to improve joint function by repeated digital manipulation of the jaws by the physiotherapist. Mobilization techniques are indicated for decreased range of motion and pain due to muscle contracture, disc displacement without reduction and fibrous adhesions of the joint. A combination of heat, cold, ultrasound and electrical stimulation is often utilized. Local anesthetic

→ DT page 8

AD

2009

Greater New York Dental Meeting


The Largest Dental Convention/Exhibition/Congress in the United States

NO Pre-Registration Fee!

MEETING DATES:
NOVEMBER 27th - DECEMBER 2nd

EXHIBIT DATES:
NOVEMBER 29th - DECEMBER 2nd

85th Annual Session



For More Information:

Greater New York Dental Meeting™

570 Seventh Avenue - Suite 800

New York, NY 10018 USA

Tel: +1 (212) 398-6922

Fax: +1 (212) 398-6934

E-mail: info@gnydm.com

Website: www.gnydm.com



Please send me more information about...

☐ Attending the Greater New York Dental Meeting

☐ Participating as a guest host and receiving free CE

☐ I speak _____ and am willing to assist international guests

Name

Address

City, State, Zip/Country Code

Telephone E-mail

Fax or mail this to:

Greater New York Dental Meeting or

visit our website: www.gnydm.com for more information.



injections are often used to improve the outcome.

Jaw Exercises

There are three types of exercises designed to achieve a different purpose. Muscle strength is addressed by isometric exercises. Isotonic exercises are used to increase range of motion. Coordination of muscle function is achieved by repetitive rhythmic exercises. Exercises should be constantly modified as symptoms change. A maintenance program is recommended with instructions on how to avoid activities that can re-injure the involved tissues and to ensure long-term resolution of symptoms.

Physical Modalities

The most common modalities used for the treatment of TMD are superficial heat and cold, ultrasound, short wave diathermy, transcutaneous electrical nerve stimulation (TENS), iontophoresis, anesthetic agents and acupuncture. The use of heat can help relax the muscle and increase blood flow to the compromised muscle. Electrotherapy devices produce thermal, histochemical and physiological changes in the muscle and joints.

Short-wave diathermy provides heat to superficial tissues whereas ultrasound can transmit heat through tissues to a depth of 5 cm.²⁶ The purpose of these modalities is to decrease pain, hyperactivity, increase tissue distensibility and may be neuromuscular re-education.^{25,26} Iontophoresis uses an electrical gradient to drive an ionic form of the medication into the tender or swollen tissues.²⁷ Acupuncture has also been used for the treatment of chronic musculoskeletal pain.²⁸

Postural Re-education

Maladaptive posture (head/neck or mandibular) may be a contributing factor in the TMD patient. The relationship between the trigeminal nerve and the upper cervical region is well recognized. Postural re-education throughout exercises and behavior modification should be considered. The participation and guidance of a physiotherapist is required for long-term stabilization of the masticatory system.

Behavioral/Psychotherapy

The TMD patient's cognitive, emotional and behavioral responses to pain are key issues in the overall evaluation and treatment. The patient's perception to pain may be maladaptive in the nature of somatization, catastrophizing and mood and subconscious habitual behavior. Failure to identify and address these factors will likely compromise the treatment outcome.

Cognitive-behavioral strategies such as behavior modification, life style counseling, progressive relaxation, guided imagery, hypnosis and biofeedback may be beneficial.^{29,30} This care is typically provided by a clinical health psychologist. Occasionally, TMD may be related to an

underlying psychosocial or psychiatric disorder such as depression or conversion disorder. In these cases, a psychiatric or clinical health psychologist referral is indicated.

Occlusal Appliance Therapy

Occlusal orthosis therapy is the most common form of treatment. They are commonly referred to as interocclusal splints, orthotic, orthosis, bite guards, bite planes, night guards and bruxism appliances. An occlusal appliance is a removable device, usually made of hard acrylic, that is custom made to fit over the occlusal surfaces of the teeth on either arch. There are generally two types of appliances: the flat plane (stabilization) appliance and the anterior reposition appliance.

The effects of appliance therapy include: prevention/reduction in abrasion to the dentition, alteration of the motor pattern of the masticatory musculature by altering periodontal ligament proprioception, alteration of muscle length, enhanced awareness of parafunctional activity, and alteration of the number, direction and quality of tooth contacts.

The major functions of non-directive, flat plane appliance therapy are muscle relaxation, dispersal of forces, enhancement of TMJ stability and the protection of teeth. Most patients are advised to utilize the appliance while sleeping or when their activity prohibits conscious awareness of daytime parafunction. Anterior repositioning appliances are used less often because repositioning of the mandible over a period of time can result in irreversible changes to the occlusion.

The purpose of these appliances is to alter the structural condyle-disc-fossa relationship in an effort to decrease adverse joint loading.^{1,31,32} Complications associated with occlusal appliance therapy arise from poor design/construction and/or excessive or incorrect use of the appliance.

Treatment of occlusion should be considered on an individual basis. Many dental conditions require treatment of the occlusion due to a lack of intra-inter-arch tooth stability, fremitus, tooth mobility, fractured teeth/restoration, and compromised function that requires redistribution of forces to minimize the effects of adverse loading.

Although dental treatment may be necessary for patients with TMD, it is believed to be very seldom necessary for the purpose of treating TMD.^{1,35} If occlusal therapy is necessary to complete treatment, it should only be initiated after the patient has regained his/her range of motion, symptoms, neuromuscular activity and when the psychosocial status is as stable as possible.

Surgery

Temporomandibular joint surgery plays a small but important role in the management of patients with TMD. The literature

shows that about 5% of the patients who undergo treatment for TMD require surgical intervention. The surgical success is dependent on a total treatment plan that involves non-surgical and surgical treatment.

Arthrocentesis is the simplest and least invasive procedure that is performed in the temporomandibular joint (TMJ). It consists of TMJ lavage placement of medications into the joint. Arthrocentesis is usually performed as an office-based procedure under local anesthesia assisted with conscious intravenous sedation. It has been suggested to be as effective as arthroscopy in the treatment of joint restriction in conditions such as internal derangement without reduction.

Surgical intervention includes arthroscopy, condylotomy, and open joint procedures, such as disk reposition and discectomy. Arthroscopy is a minimally invasive technique that is usually performed under general anesthesia. It is primarily performed in the upper joint space, and it's useful for minor debridement and lavage, incision of minor adhesions and biopsies.

Indications for surgery of the TMJ may be divided into relative and absolute. Absolute indications are reserved for conditions such as tumors, growth abnormalities and ankylosis of the TMJ. The general indications for TMJ surgery for the most common disorders, internal derangement and osteoarthritis, are significant TMJ pain and dysfunction that are refractory to nonsurgical treatment, and there is imaging evidence of disease. It has been suggested that arthrocentesis and arthroscopy be used for TMJ pain and limited opening, and open TMJ surgery should be reserved for advanced cases of TMJ derangement and osteoarthritis.³⁴

Based on our current understanding of TMD, successful management is dependent upon recognizing the complexity of this condition in terms of the multifactorial nature and the need in most cases of a multidisciplinary approach. Recognizing that on many occasions these musculoskeletal conditions may have a co-existing psychosocial component that may need to be addressed in order to enhance the treatment outcome is also important. A complete and accurate diagnosis on a case-specific basis will provide the development of the most efficacious individualized approach to care. DT

Literature

1. Okeson, J.P., ed. Orofacial pain: guidelines for assessment, diagnosis and management. Chicago: Quintessence Publishing Co., Inc., 1996: 113-84.
2. Mejersjo, C., and Carlsson, G.E. Long-term results of treatment of temporomandibular pain dysfunction. J Prosthet Dent, 1983, 49:809-15.
3. Okeson, J.P. Management of Temporomandibular Disorders and Occlusion, 2nd ed. St Louis: CV Mosby Co., 1989.
4. Nickerson, J.W., and Boering, G. Natural course of Osteoarthritis as it relates to internal derangement of the temporomandibular joint. Oral and Maxillofac Surg Clin North Am, 1989, 1:27-46.

5. Greene, C.S., and Marbach, J.J. Epidemiologic studies of mandibular dysfunction. A critical review. J Prosthet Dent, 1982, 48:184-90.
6. Carlsson, G.E. Long-term effects of treatment of craniomandibular disorders. J Craniomand Pract, 1985: 3:337-342.
7. Brown, D.T., Gaudet, E.L., Outcome measurement for treated and untreated TMD patients using the TMJ scale. J Craniomand Pract 1994; 12:216-221.
8. Garafis, P. Grigoriadou, E., Zarafi A., et al. Effectiveness of conservative treatment for craniomandibular disorders: A 2-year longitudinal study. J Orofacial Pain 1994; 8:509-514.
9. Rasmussen, O.C. Clinical finding during the course of temporomandibular arthropathy. Scan J Dent Res 1981; 89:283-288.
10. Helkimo, E., Westling, L. History, clinical findings, and outcome of treatment of patients with anterior disc displacement. J Craniomand Pract 1987; 5:270-276.
11. Rasmussen, O.C. Temporomandibular arthropathy. Int J Oral Surg 1983; 12:365.
12. Rugh, J.D., and Harlan, J. Nocturnal bruxism and temporomandibular disorders. Adv Neurol 1988; 49:329-41.
13. Rugh, J.D., and Johnson, R.W. Temporal analysis of nocturnal bruxism. J Periodontol 1981; 52:263-65.
14. Gangarosa, L.P., Mahan, P.E. Pharmacologic management of TMD-MPDS. Ear Nose Throat J 1982; 61:670-678.
15. Gregg, J.M., Rugh, J.D., Pharmacological therapy. In: Mohl, N.D., Zarb, G.A., Carlsson, G.E., et al. (eds). A Textbook on Occlusion. Chicago: Quintessence 1988: 351-375.
16. Wenneberg, B., Kopp, S. Short term effects of intra-articular sodium hyaluronate, glucocorticoids, saline injection on rheumatoid arthritis of the temporomandibular joint. J Craniomandib Disord Facial Oral Pain 1991; 5:251-258.
17. Wenneberg, B., Kopp, S. Grondahl, H.G. Long term effect of intra-articular injections of a glucocorticosteroid into the TMJ: A Clinical and radiographic 8 year follow up. J Craniomandib Disord Facial Oral Pain 1991; 5:11-18.
18. Dellemijn, P.L.L., Fields, H.L. Do benzodiazepines have a role in chronic pain management? Pain 1994; 137-152.
19. Harkins, S., Linford, J., Cohen, J., et al. Administration of Clonazepam in the treatment of TMD and associated myofascial pain: a double-blind study. J Craniomandib Disord Facial Oral Pain 1991; 5:179-186.
20. Stanko, J.R. A review of oral skeletal muscle relaxants for the craniomandibular disorder (CMD) practitioner. J Craniomand Pract 1990; 8:234-243.
21. Gangarosa, L.P., Mahan, P.E. Pharmacological management of TMD-MPDS. Ear Nose Throat J 1982; 61:30-41.
Brown, R.S., and Bottomley, W.K. The utilization and mechanisms of tricyclic antidepressants in the treatment of chronic facial pain: A review of the literature. Anesth Prog 1990; 37:223-9.
22. Danzig, W., VanDyke, A.R. Physical therapy as an adjunct to temporomandibular joint therapy. J Prosthet Dent 1983; 49:96-99
23. Clark, G.T., Adashi, N.Y., Dornan, M.R. Physical medicine procedures affect temporomandibular disorders: A review. J Am Dent Assoc 1990; 121:151-161.
24. Carlson, C.R., Okeson, J.P., Falace, D.A., et al. Stretch based relaxation and the reduction of EMG activity among masticatory muscle pain patients. J Craniomandib Disord Facial Oral Pain 1991; 5:205-212.
25. Au, A.R., Klineberg, I.J. Isokinetic exercise management of temporomandibular joint clicking in young adults. J Prosthet Dent 1993; 70:33-39.
26. Mohl, N.D., Ohrback, R.K., Crowe, H.C., Gross, A.J. Devices for the di-

agnosis and treatment of temporomandibular disorders, Part III. Thermography, ultrasound, electrical stimulation and EMG biofeedback. J Prosthet Dent 1990; 63:472-7.
27. Lark, M.R., Gangaros, L.P. Iontophoresis: An effective modality for the treatment of inflammatory disorders of the temporomandibular joint and myofascial pain. J Craniomand Pract 1990; 8:108-119.
28. Johansson, A., Wenneberg, B., Wagersten, C., et al. Acupuncture in the treatment of facial muscular pain. Acta Odontol Scand 1991; 49:153-158.
29. Rugh, J.P. Psychological components of pain. Dent Clin North Am 1987; 31:579-94.
30. Moss, R.A., Adams, H.E. The assessment of personality, anxiety and depression in mandibular pain dysfunction subjects. J Oral Rehabil 1984; 11:233-7.
31. Clark, G.T. A critical evaluation of orthopedic interocclusal appliance therapy: Design, theory and overall effectiveness. JADA 1984; 108: 359-64.
32. Maloney, F., Howard, J.A. Internal derangements of the temporomandibular joint. III. Anterior repositioning splint therapy Aus Dent J 1986; 31:130-9.
33. McNeill, C. editor. Craniomandibular disorders: guidelines for evaluation, diagnostic and management. Chicago: Quintessence Publishing Co Inc, 1990.
34. Dolwick, M.F. Temporomandibular Joint Surgery for internal Derangement. Dent Clin N Am 51:2007; 195-208.

MEDIA CME

Self-Instruction Program

Dental Tribune Middle East & Africa in collaboration with CAPP introduce to the market the new project mCME - Self Instruction Program.

mCME gives you the opportunity to have a quick and easy way to meet your continuing education needs.

mCME offers you the flexibility to work at your own pace through the material from any location at any time. The content is international, drawn from the upper echelon of dental medicine, but also presents a regional outlook in terms of perspective and subject matter.

How can professionals enroll? They can either sign up for a one-year (10 exercises) by subscription for the magazine for one year (\$65) or pay (\$20) per article. After the payment, participants will receive their **membership number** and will be able to attend to the program.

How to earn CME credits? Once the reader attends the distance-learning program, he/she can earn credits in three easy steps:

1. Read the articles.

2. Take the exercises

3. Fill in the Questionnaire and Submit the answers by Fax (+971 4 3686885) or

Email :info@cappmea.com

After submission of the answers, (name and membership number must be included for processing) they will receive the Certificate with unique ID Number within 48 to 72 hours.

Articles and Questionnaires will be available in the website after the publication.

www.cappmea.com

We are currently undertaking more revolutionary projects

An interview with Mectron, the company who invented Piezosurgery



Fernando Bianchetti



Domenico Vercellotti



Wolf Narjes



Alexandre Cadau

Mectron, based in Italy, has revolutionised dental surgery with their development of piezoelectric bone surgery. Recently, the company presented the 3rd generation of their Piezosurgery device at the IDS show in Cologne in Germany. We spoke with company founders Domenico Vercellotti and Fernando Bianchetti, as well as area managers Wolf Narjes and Alexandre Cadau, about the clinical advantages of their invention and how the company is reacting to the current market conditions.

Dental Tribune: Market prospects for 2009 are rather uncertain due to the financial crisis. Is your company prepared for a potential economic slowdown?

Fernando Bianchetti: The only way to withstand this crisis is to remain successfully in the market through investments in scientific and technical research, in Europe and other countries.

Domenico Vercellotti: What Fernando just said has always been our corporate philosophy; it will certainly help us in difficult times like this. Mectron offers high quality products at reasonable prices and puts a lot of effort into the development of new technologies and not merely into expensive marketing campaigns.

Wolf Narjes: Being a family-owned company, Mectron is probably more flexible and manageable than larger companies. Therefore, we can react relatively quickly to unexpected market changes.

Have you already experienced an economic climate change in Italy and other markets?

Fernando Bianchetti: Since our company was founded in

1979, we have already had to go through occasional tough economic times. However, nothing really compares with the latest financial crisis.

been studied, to ensure that there is no risk for users and patients and that the medical effects are always positive. Many companies have at-

Alexandre Cadau: Piezosurgery has certainly been one of the most important developments in the dental and medical field. This unique device

the whole clinical research and training activities in Piezoelectric Bone Surgery and works independently from Mectron.

Wolf Narjes: It is essential to be suitably trained in this technique. Therefore, we offer courses in Europe, Asia, as well as North and South America. Last year, we opened a new branch in Phuket in Thailand that serves as the Piezosurgery training centre for the whole Asia Pacific Region.

Alexandre Cadau: There is a reason that training is crucial for Piezosurgery. Users experience a steep learning curve before getting used to the micrometric movement of Piezosurgery, which is completely different to the traditional techniques. We organise workshops in many countries around the world that help dentists learn the differences between Piezosurgery and conventional burs and saws. In addition, we collaborate with universities, to offer attending clinicians cadaver dissection courses that help them appreciate the surgical benefits.

“Piezosurgery has certainly been one of the most important developments in the dental and medical field.”

Alexandre Cadau: Fernando is right. At the moment, we are experiencing a huge loss of confidence in all consumer groups. On the other hand, we have always been challenged by the depreciation of various foreign currencies, like in 1992 when devaluation hit many countries.

Your company is mainly known for its innovative Piezosurgery technology. What are the main advantages compared to traditional surgical technologies?

Domenico Vercellotti: Mectron invented piezoelectric bone surgery in collaboration with Prof. Tomaso Vercellotti almost ten years ago. Back then, it was not just another product: it was a significant innovation in the field of dentistry based on technical expertise and years of clinical research. Thanks to Piezosurgery, oral surgery evolved from traditional rotating instruments to a new system of cutting bone that spares soft tissue and accelerates the healing process.

Wolf Narjes: Our Piezosurgery device is scientifically approved and we are considered to be the only company in this field to have a clinical database on each available surgical instrument. All the clinical applications for the device have

tempted launching similar products, but they are still missing scientific data or research regarding the effectiveness of their methods.

Would you tell us more about how Piezosurgery was developed and how it has been received in different markets?

Wolf Narjes: I have found that several countries, including South Korea, Italy, and Germany, have been very open-minded to this new technology. Most Scandinavian countries, however, have only begun to understand how to use this innovative technique.

allows the surgeon to work in less stressful and safer conditions. Postoperative healing times are also reduced three-fold with this method.

Domenico Vercellotti: The latest innovation is tips for the implant site preparation that have demonstrated histological benefits and a better osseointegration of implants compared to the traditional twist drill (Giulio Preti et al., ‘Cytokines and Growth Factors Involved in the Osseointegration of Oral Titanium Implants Positioned using Piezoelectric Bone Surgery Versus a Drill Technique: A Pilot Study in

“It is essential to be suitably trained in this technique.”

Fernando Bianchetti: All the clinical protocols and techniques developed for Piezosurgery are based on scientific publications endorsed by universities and credible specialists in the field of dental surgery. They confirm not only the benefits for the clinician, such as maximum surgical precision and wider intra-operative visibility, but also those for patients who suffer from less postoperative pain.

Minipigs’, *Journal of Periodontology*, 78 (2007): 716-722).

Training courses are regularly offered at the Piezosurgery Academy in Italy. Do you also offer courses in other parts of the world?

Domenico Vercellotti: Piezosurgery Academy was established by Prof. Tomaso Vercellotti to give scientific support to the Piezoelectric Bone Surgery. It is managing

With four regional headquarters, do you consider yourself a global cooperation?

Fernando Bianchetti: Certainly, our branches in Germany, India, and the Asia Pacific region report to our headquarters in Italy. In other countries, we have worked successfully with local dealers for almost ten years, in some countries even 20 years.