


implants

international magazine of oral implantology



research

The dos and don'ts in the handling of PRF

case report

Peri-implant bone regeneration
through laser decontamination

industry

The next generation biomaterial
for soft-tissue augmentation



IT'S SIMPLE TO BE A
WINNER



**MIS GLOBAL
CONFERENCE**

May 14-17, 2020
Marrakech
Morocco



**PROVEN SUCCESS MEETS ENHANCED
STABILITY. MAKE IT SIMPLE**

The biological stability and predictable esthetics of the SEVEN, combined with the extensive research and development process have given the SEVEN a potential advantage in soft tissue preservation and growth as well as an array of restorative benefits. Learn more about the SEVEN implant system and MIS at: www.mis-implants.com

**MIS[®]
SEVEN[®]**



Dr Rolf Vollmer

First Vice President and Treasurer of DGZI



Join **DGZI** on the track for success

With a 50-year history, the German Association of Dental Implantology (DGZI) represents a practice-oriented and evidence-based approach to implantology in Germany. DGZI always keeps pushing boundaries in this particularly innovative field of dentistry. We place an overarching focus on actively supporting our practising colleagues and dental technicians by offering a wide range of special training courses. I can proudly say that DGZI is one of the best internationally networked German expert societies for dental implantology, boasting more than registered 4,000 members in Germany alone and more than 13,000 cooperating members in 66 countries abroad. In addition, the annual meeting of DGZI is considered a definite highlight and it is fair to say that it has become an integral part of the annual schedules of numerous clinicians from all around the globe.

With more than 300 holders of the DGZI specialist in oral implantology certification, our expert society can recommend a vast number of highly qualified clinicians who continue to break the mould every single day in their dental practices. Thanks to our close ties to universities and research facilities we can implement the latest findings effortlessly into their daily practice. Fully recognised by the "Konsensuskonferenz Implantologie" (a joint initiative of German professional associations involved in dental implantology), the one-year DGZI implantology curriculum is the base for the practitioner. It features an innovative e-learning concept, which consists of regularly updated compulsory and optional modules, and is now offered in English too. The DGZI implantology curriculum is a state-of-the-art beacon of education for young aspiring dentists looking to delve deeply into the specialist field that is oral implantology.

In 2019 the American Board of Oral Implantology (ABOI) in the US decided to make the ABOI/ID Diplomate examination available for experienced dental practitioners internationally. The ABOI has an independent examination committee chartered by the American Academy of Implant Dentistry (AAID), the official US partner of DGZI. Now, especially graduates of the DGZI implantology curriculum, as well as holders of both Expert in Dental Implantology and DGZI Specialist in Oral Implantology certification can take this examination to become an ABOI/ID Diplomate in addition to the already gained credentials. DGZI can offer individual training and/or preparatory seminars for those wishing to pursue this prestigious certification on request.

Last but not least we should not forget our endeavours in publishing: The German language *Implantologie Journal* and the English language *implants—international magazine of oral implantology*. The former is DGZI's personal member journal and delivered to our German DGZI members, oral surgeons, prosthodontists and dental technicians on a monthly basis. The latter, with a circulation of 10,000 copies and four issues every year, is popular in over 90 countries around the world.

I wish you a great and interesting time reading this first 2020 issue of *implants—international magazine of oral implantology*.

Yours,

A handwritten signature in black ink, appearing to read 'R. Vollmer'.

Dr Rolf Vollmer



page 06



page 16



page 32

Cover image courtesy of
TBR Dental Group
www.tbr.dental



editorial

- Join **DGZI** on the track for success 03
Dr Rolf Vollmer

research

- The dos and don'ts in the handling of **PRF** 06
Prof. Shahram Ghanaati, Dr Sarah Al-Maawi, Dr Eva Dohle & Dr Torsten S. Conrad
- Success rate of dental implants in **heavy smokers** 12
Drs Branislav Fatori & Inge Schmitz

case report

- Implant-retained **overdenture** on a very **thin** bone ridge 16
Drs Nicola Alberto Valente, Murali Srinivasan & Nicole Kalberer
- Peri-implant** bone regeneration through **laser** decontamination 22
Prof. Wilfried Engelke, Dr Christian Engelke, Dr Victor Beltrán & Dr Marcio Lazzarini
- Sixteen years of **follow-up** after insertion of a Z1 implant 28
Dr Virgilio Masini

industry

- The **next generation** biomaterial for soft-tissue augmentation 32

practice management

- Feedback done **right** 34
Andrea Stix, MSc, MBA

interview

- Reshaping** implantology 36
- The **link** between oral disease and **oxidative stress** 38

news

- manufacturer news 40
- news 44

about the publisher

- imprint 50

TRULY CONICAL PRECISION

CONELOG[®]
SYSTEM



CONELOG[®] PROGRESSIVE-LINE conical performance^{1,2} at bone level

Precise conical connection

- long conus for reduced micromovements
- superior positional stability in comparison to other conical systems^{1,2}
- easy positioning with excellent tactile feedback
- integrated platform switching supporting the preservation of crestal bone

[1] Semper-Hogg, W, Kraft, S, Stiller, S et al. Analytical and experimental position stability of the abutment in different dental implant systems with a conical implant-abutment connection Clin Oral Invest (2013) 17: 1017

[2] Semper Hogg W, Zulauf K, Mehrhof J, Nelson K. The influence of torque tightening on the position stability of the abutment in conical implant-abutment connections. Int J Prosthodont 2015;28:538-41



The dos and don'ts in the handling of PRF

Prof. Shahram Ghanaati, Dr Sarah Al-Maawi, Dr Eva Dohle & Dr Torsten S. Conrad, Germany




Fig. 1: Visualisation of the puncture direction during blood collection.

Autologous blood concentrates, and platelet-rich fibrin (PRF) in particular, are increasingly used today to support wound healing and regenerative processes.¹ PRF is made from the patient's own peripheral blood without the addition of anticoagulants. A solid or liquid PRF matrix can be obtained through a single centrifugation process, depending on the collection tube that is used.² Through this centrifugation process, the blood components are separated according to the centrifugal force used. The red blood cells move towards the bottom of the tube.³ The platelets and leucocytes are concentrated in the upper layer, the remaining fibrin matrix. Thus, this autologous blood concentrate, which also contains further plasma proteins, is capable of actively releasing different growth factors such as vascular endothelial growth factors (VEGF), epidermal growth factors (EGF), or platelet-derived growth factors (PDGF) over a relatively long period of time (up to fifteen days).^{4,5} These growth factors play a key role in the support of wound healing and regenerative processes, since they contribute to the formation of new vessels, epithelialisation and the stimulation of further regener-

ative cells.^{6,7} The composition and bioactivity of PRF depends primarily on the centrifugal force that is used during centrifugation.³

Several recent studies have demonstrated the influence of the centrifugal force on the composition and bioactivity of the obtained PRF.⁸⁻¹² It has been shown that the application of a low centrifugal force for accumulation leads to a significantly higher number of platelets and leukocytes in PRF compared to a medium or high centrifugal force.^{3,10} Growth factors are released in a similar way. PRF matrices that are prepared with a low centrifugal force release significantly higher concentrations of different growth factors (such as VEGF, PDGF, EGF, TGF- β 1) compared to PRF matrices that are prepared using a higher centrifugal force.⁸⁻¹² As a result, the so-called Low-Speed Centrifugation Concept (LSCC) was introduced, which aims at standardising the production of blood concentrates and enabling reproducible treatment protocols or clinical results.³ This article will particularly focus on the technical aspects of the clinical application and handling of PRF. The tubes used for the production of PRF have been specifically developed for this particular purpose. Depending on the clinical indication, two different variants of PRF matrices exist. PRF tubes with a glass surface promote coagulation. During centrifugation, a solid PRF matrix is formed. In contrast, the coagulation process can be slowed down by means of plastic-coated tubes. Accordingly, coagulation is slowed down during centrifugation. At room temperature, a PRF matrix remains liquid for about thirty minutes after centrifugation until it eventually coagulates.

Protocol	RPM (x 100)	Duration (min)	Centrifugal force (x G)
High concentration RCF	2,400	8	710
Medium concentration RCF	1,200	8	177
Low concentration RCF	600	8	44

RCF, relative centrifugal force; RPM, revolutions of the centrifuge per minute.

Table 1: Visualisation of the different LSCC protocols (Low-Speed Centrifugation Concept for a centrifuge with a radius of 110 mm).

Blood collection

For the production of PRF the patient's own venous blood is required, which is taken from the peripheral veins after the patient has been fully briefed on the procedure. This blood collection is a routine method and is particularly used in diagnostics. The blood collection should be carried out according to the guidelines of the World Health Organization (WHO).¹³ In order to find a suitable puncture site, the anatomical position of the peripheral veins should first be palpated. For this purpose, the vena mediana cubiti, which is located in the antecubital fossa (inner bend of the elbow), is ideally suited. Gloves must be worn and the tourniquet must be placed approximately 5 cm above the puncture site of the vein, which must be disinfected with a skin antiseptic according to the manufacturer's instructions. A butterfly needle is then inserted into the vein at an angle of 30° to the skin surface (Fig. 1). To avoid completely piercing through the vein, the angle should be

flattened once the vein has been hit. The vacuum system of the PRF tube then fills the tube with venous blood until an amount of 10 ml is reached and the blood supply can be stopped automatically. After loosening the tourniquet tube, the butterfly cannula can be removed. Subsequently, sufficient pressure should be exerted on the puncture site with a sterile swab in order to avoid secondary bleeding underneath the skin.

Centrifugation

In order to avoid early physiological coagulation of the blood, the PRF tubes must be centrifuged quickly after blood collection in a dedicated centrifuge that stands on a table nearby in the same treatment room. Through centrifugation, a separation process is triggered, which sediments cells and/or biomolecules from a suspension (i.e. blood), depending on the relative centrifugal force and the size, shape and density of the vari-




Fig. 2: Balance pattern of the centrifuge when loading two, three, four and six tubes. Centrifugation of five, seven, nine or eleven tubes is not possible. For this purpose, an additional tube filled with water must be used.




Fig. 3: Separating the red phase from the solid PRF phase. **Fig. 4:** A PRF plug. **Fig. 5:** A thinly pressed PRF matrix.

ous suspension components. The relative centrifugal force (RCF) represents the centrifugal force as a multiple of the Earth's gravity and is often expressed as the value G. Moreover, it is a decisive factor for the resulting concentration of the sedimented cells and biomolecules in PRF. The value G indicates exactly the force required for an optimal centrifugation of a corresponding suspension (in this case blood) to obtain the desired sediment (in this case PRF) as final product, and allows the calculation of the necessary speed of the centrifuge rotor for a corresponding tube and centrifuge.

For centrifuges where only the rotational speed (revolutions of the centrifuge per minute, RPM) can be set, the RCF or the necessary G-value must first be calculated by means of a fixed formula. The relation between rotational speed (RPM) and RCF depends on the size of the rotor (r = radius of rotation = distance between the axis of rotation and the bottom of the tube). Here, the following formula is used for conversion:¹⁴ $RCF = 1.12 \times \text{radius} \times (\text{RPM}/1,000)^2$. The relative centrifugal force required for PRF production using the established LSCC

is given in Table 1 and should be set on the centrifuge according to the clinical indication ($r = 110\text{mm}$). In general, the centrifuge should be placed on a stable and even fundament. When loading the centrifuge with the blood-filled tubes, it is imperative to ensure that any imbalance is eliminated. This means that the tubes must be placed inside of the rotor in such a way that the weight of the tube placed exactly opposite the other is identical (Fig. 2). If the number of tubes is uneven, a tube filled with the equivalent volume (such as sodium chloride, for example) must be added to compensate for the weight.

Processing of PRF

Immediately after centrifugation, the tubes are carefully removed from the centrifuge and transferred to an appropriate tube holder. Owing to the applied RCF and depending on the size, shape and density of the blood components, only two phases can now be visually identified: a red phase at the bottom of the tube, which contains mainly erythrocytes and a PRF phase on top of it, filling up the upper part of the tube. In the case of solid PRF, which is obtained by centrifuging the blood in the red PRF tubes, these two phases coagulate very quickly. In order to separate the solid PRF matrix from the red lower phase, it is recommended to first separate the two phases roughly by cutting them with scissors. In concrete terms, this means carefully lifting up the upper phase of solid PRF with sterile tweezers (the lower red phase is lifted as well) and then roughly separate the two phases in the upper part of the red phase (Fig. 3). The PRF phase (with remains of the red phase) is then transferred into a dedicated PRF box provided for this particular purpose. This PRF box, which has been specially developed for various PRF indications, consists of a stainless-steel container with a self-weighted lid and a large and small stamp. In this box, the remaining parts of the red phase can now be removed from the PRF phase by carefully wiping it off with a blunt object (such as a closed pair of scissors). Thereafter, solid PRF matrices can be further




Fig. 6: Separating liquid PRF from the red phase by means of a syringe.

Conical? Parallel?

copaSKY!

The unique hybrid connection
for sophisticated restorations!



Leading in Immediate Restoration – powered by physiological Prosthetic!

DENTAL INNOVATIONS
SINCE 1974

bre^{dent} group

bre^{dent} UK Ltd · Unit 26, Tapton Park Innovation Centre · Brimington Road · Chesterfield S41 0TZ · T: +44 1246 559599 · F: +44 1246 557777 · www.bredent.co.uk · @: enquiries@bredent.co.uk