

laser

international magazine of laser dentistry



ISLD



research

Diverse applications of lasers in dentistry

case report

Diode laser-assisted vital pulp therapy

events

1st IDSCLD in Plovdiv, Bulgaria





16th WORLD CONGRESS WFLD 2018

1 – 3 OCTOBER 2018
RWTH AACHEN UNIVERSITY
AACHEN, GERMANY

**RWTHAACHEN
UNIVERSITY**



The 16th Congress of the
World Federation for Laser
Dentistry (WFLD)

The 27th Annual Meeting of
the German Society for Laser
Dentistry (DGL)

The 6th Annual Congress
of the World Academy for
Laser Education
in Dentistry (WALED)

This congress will integrate science and practical experience on different levels of presentation and demonstrations, like:

- High ranked international keynote speaker lectures
- On stage live patient demonstrations
- Interactive digital poster presentations
- Oral presentations combined with clinical relevant skill training
- Short presentations of latest research findings
- Clinical case presentations
- Rotating company supporting workshops, gaining continuous education certificates.

CONTACT

Headquarters WFLD 2018 Aachen
Mr. Leon Vanweersch
Vice-Organizing Chairman
Phone: +49 151 50610781
headquarters@wfl-d-aachen2018.com
www.wfl-d-aachen2018.com

Prof. Dr Norbert Gutknecht

DGL President & Organising Chairman
of the 16th WFLD World Congress



Anniversary congress in Aachen from 1 to 3 October 2018

Dear friends of laser dentistry,

It is my great pleasure to be able to welcome you in Aachen, Germany, for this extraordinary congress event: 30 years of WFLD/ISLD, 27 years of DGL and 12 years of WALED. With this anniversary congress, we are making it possible for many colleagues from all over the world to jointly participate in the scientific presentations and clinical treatment sessions.

If we are considering the application of lasers in dentistry, we have to realise that while it is a well-established treatment method today, it was only initiated 30 years ago. Diverse universities have conducted research which in the end has led to a scientifically ensured application of these methods. At the same time, scientists and doctors have come together in working groups which have eventually become the foundation for the establishment of laser societies. Today, nobody would question an indication-based application of lasers in dentistry anymore.

The ISLD (International Society for Laser Dentistry), as well as the DGL were founded with the objective to promote the application of lasers in dentistry in both science and practice. In 1998, the DGL and ISLD thus became associated and have been closely collaborating ever since.

As part of this collaboration, the DGL successfully organised the ISLD World Congress in Berlin in 2006. As one of the founding members of DGL and Organising Chairman, I am especially happy and honoured that

the DGL will again be the hosting society of the anniversary congress of WFLD/ISLD in Aachen, Germany, and that Prof. Dr Lynn Powell as a founding member of ISLD has taken over the position as Chairman of the Scientific Committee.

We will offer scientists and practitioners the scientific and clinical platform to present and discuss their research results and clinical cases, and to stay up-to-date with the developments and possibilities, as well as with the limits of applying lasers in oral medicine. The workshops offered as part of the congress are dedicated to continuous clinical education and are thus acknowledged with further education credits.

As you know, it is not only the aim of a congress to further educate oneself scientifically and clinically but also to socialise with friends and colleagues. I am thus looking forward to enjoying good food and live music with you at Rahe Castle on Tuesday evening.

With all this in mind, I would like to wish us all a harmonious congress.

Yours

Prof. Dr Norbert Gutknecht



Cover image courtesy of Light Instruments Ltd., in cooperation with Swarovski www.light-inst.com



editorial

Anniversary congress in Aachen from 1 to 3 October 2018 03
Prof. Dr Norbert Gutknecht

research

Diverse **applications** of **lasers** in dentistry 06
Dr Igor Cernavin

case report

Diode laser-assisted **vital pulp therapy** in pulp polyp treatment 10
Drs Maziar Mir, Masoud Mojahedi, Jan Tunér & Masoud Shabani

industry

Dual-wavelength laser in **aesthetic** and **prosthetic** dentistry 14
Dr Bogdan Crisan *et al.*

practice management

Six Sigma in the management of **laser-assisted** dental practices 20
Dr Imneet Madan, UAE

Successful **communication** in your **daily practice** 24
Dr Anna Maria Yiannikos

interview

Launching a **novel cutting-edge** dental laser line 26

events

1st International Dental Students Congress for Laser Dentistry 30
Dr Georgi Tomov

news

manufacturer **news** 28

news international 32

DGL

Jubiläumskongress vom 1. bis 3. Oktober 2018 in **Aachen** 37
Prof. Dr. Norbert Gutknecht

news germany 38

about the publisher

imprint 42

Fotona
choose perfection

The universe at your fingertips.

● LightWalker

The highest technology dental laser system

Supreme clinical results:


- TwinLight® Perio Treatments
- TwinLight® Endo Treatments
- No-sutures soft-tissue surgery
- Patient-friendly conservative dentistry
- Pre-sets for over 40 applications


Unmatched simplicity of use:

- Balanced and weightless OPTOflex® arm
- Nd:YAG handpiece detection system
- Quantum Square Pulse technology for fast minimally invasive treatments
- X-Runner™ - the first digitally controlled Er:YAG dental laser handpiece

Journey into a new dental experience with speed, precision and great results.

Visit www.fotona.com today!



 Fotona App





© tandaV/Shutterstock.com

Diverse applications of lasers in dentistry

Recent literature

Dr Igor Cernavin, Australia

When considering whether to work with lasers and in which field they could be applied, recent studies provide many application options and issues for practitioners to consider. The following presents some of the newest research on possible areas of application and further investigation.

Petrov et al. used a femtosecond laser with a high repetition rate, which is probably the future of lasers for hard-tissue removal to achieve fast and more precise ablation in dentine and enamel.¹ They concluded that the ultra-fast femtosecond laser used in their work holds the promise of a significant drilling ability without collateral thermomechanical effects. It achieved high processing efficiency, overcame disadvantages of other laser sys-

tems reported, and can be used to develop an instrument for cavity preparation based on fast and precise ablation. Their further aim is to exceed the speed of conventional drilling instruments and thus to reduce the treatment time, which in turn will bring comfort to the patient.

Levine published an article on how to choose the right laser for one's practice, which readers may find of interest.²

Hashimoto et al. investigated fluoridated hydroxyapatite for application as an implant coating for titanium bone substitute materials for dental implants.³ They concluded that fluoridated hydroxyapatite coatings are suitable for real-world implantation applications.

Giannelli et al. carried out a double-blind, randomised, single-centre, split-mouth clinical trial investigating the efficacy of and patient-reported outcomes after one year of treatment of severe periodontitis with a laser and light-emitting diode (LED) procedure adjunctive to scaling and root planing.⁴ Their study confirmed the efficacy of combined phototherapy and scaling and root planing, which had emerged from previous clinical trials, extending its field of application to severe periodontitis.⁴

Belcheva et al. carried out a study whose aim was to evaluate the positive effects of the carbon dioxide laser (10,600nm) with acidulated phosphate fluoride gel on enamel acid resistance.⁵ Their conclusion was that this combination was more effective in protecting the enamel surface and resisting demineralisation than was carbon dioxide laser irradiation or fluoride alone.⁵

Campos et al. published a double-blind study on immediate laser-induced haemostasis in anticoagulated rats subjected to oral soft-tissue surgery.⁶ There has been much controversy about the management of patients on oral anticoagulants requiring oral surgical procedures. The haemostatic properties of high-power lasers were perceived to be potentially helpful during oral soft-tissue surgeries in anticoagulated patients. The authors concluded that laser-induced haemostasis is an alternative for intra- and postoperative bleeding control in patients on anticoagulation therapy.⁶

As oncological treatment can result in changes in the oral cavity, Carvalho et al. drafted a guide, based on a systematic review, directed at the team of health professionals involved in the oral care of oncological patients.⁷ The review concentrated on randomised clinical trials involving paediatric and adult oncological patients, focusing on the prevention and treatment of oral complications.⁷ The studies included in the review emphasise the provision of Low Level Laser Therapy, among other interventions, to minimise the severity of oral problems in such patients.⁷

Tani et al. carried out an *in vitro* study that compared photo-biomodulation potentiality using red (635 ± 5 nm) or near-infrared (808 ± 10 nm) diode lasers and violet-blue (405 ± 5 nm) LED operating in a continuous wave with a 0.4 J/cm energy density, on human osteoblast and mesenchymal stromal cell viability, proliferation, adhesion and osteogenic differentiation.⁸ They concluded that the 635 nm laser had a potential effective option for promoting/improving bone regeneration.⁸

Ghouth et al. carried out a systematic review of the evidence on the use of laser Doppler flowmetry in the assessment of the pulpal status of permanent teeth compared with other sensibility and/or vitality tests. They con-

cluded that, despite the higher reported sensitivity and specificity of laser Doppler flowmetry in assessing pulp blood flow, this data is based on studies with a high level of bias and serious shortfalls in study design.⁹ More research is needed to study the effect of different laser Doppler flowmetry's parameters on its diagnostic accuracy and the true cut-off ratios by which a tooth could be diagnosed as having a normal pulp.⁹

Kaur et al. compared soft-tissue wound healing using diode lasers (810nm) versus the conventional scalpel approach as an uncovering technique during second-stage surgery for implants.¹⁰ They found that it can minimise surgical trauma, reduce the amount of anaesthetic required, improve visibility during surgery owing to the absence of bleeding and eliminate postoperative discomfort.¹⁰

Efficiency in debonding porcelain laminate veneers was studied by Al-Balkhi et al. using several laser parameters and two different application modes of the Er:YAG laser (contact and non-contact mode).¹¹ Their finding was that the Er:YAG laser is an effective tool in debonding porcelain laminate veneers. The non-contact application mode was more efficient in reducing debonding time than the contact application mode, but resulted in a higher change in pulp temperature.¹¹

Kellesarian et al. carried out a comprehensive review to assess the effectiveness of erbium lasers in the removal of all-ceramic fixed dental prostheses and found that the benefits of lasers over mechanical instrumentation for crown removal encompassed efficient restoration retrievability without restoration or tooth surface damage and a relatively easier and more time-effective procedure with no prerequisite for anaesthetic agents.¹² It is, however, imperative for clinicians to be well trained and exhibit adequate knowledge regarding recommended power settings and laser-safety parameters with reference to interactions between light and different tissues and ceramics.¹²

The effect of Er:YAG (Smart 2940D Plus, DEKA) and Er,Cr:YSGG (Waterlase iPlus, BIOLASE) lasers on the shear bond strength between orthodontic brackets and dental porcelain in comparison

© BarGar/Shutterstock.com

with conventional acid etching with 9% hydrofluoric acid (Ultradent Products) was investigated by Mirhashemi et al.¹³ They concluded that with the laser groups the failures were mostly adhesive, while they were mostly cohesive with the controls.¹³ They found that the Er:YAG laser with the specifications they used was not a suitable alternative to hydrofluoric acid etching.¹³ In the case of the Er,Cr:YSGG laser, although the conditioning outcome met the bond strength requirement for orthodontic brackets (6–8MPa) they concluded that the bond strength must be further improved by fine-tuning the irradiation parameters.

Yassaei et al. assessed the efficacy of an Er:YAG laser and pastes containing casein phosphopeptide-amorphous calcium phosphate (CPP-ACP) with and without fluoride and their combination for prevention of white spot lesions in the enamel.¹⁴ They found that the Er:YAG laser was able to decrease demineralisation.¹⁴ It further proved to be a potential alternative to preventative dentistry and was more effective when combined with CPP-ACP products.¹⁴ This would be useful especially for orthodontics.

Sarmadi et al. evaluated patients' experiences of two excavation methods, the Er:YAG laser and rotary bur, and the time required with these methods, as well as objective assessments of quality and durability of restorations over a two-year period.¹⁵ Their conclusions were that the Er:YAG laser technique was more time-consuming than the rotary bur, but despite this, the laser technique caused less discomfort and was preferred as an excavation method by patients.¹⁵

Li et al. carried out a meta-analysis to systematically evaluate the applications of Er:YAG lasers for the removal of caries and cavity preparation in children.¹⁶ They concluded that the time required for Er:YAG laser treatment was longer than that for the conventional mechanical method, but there was less pain associated with the Er:YAG laser treatment.¹⁶ There were no significant differ-

ences in the complete retention rate, marginal discoloration and marginal adaptation when compared with the conventional method.¹⁶

Pinheiro et al. assessed the utility of dental acid etchants containing 37% phosphoric acid and methylene blue dye as a sensitising agent for photodynamic therapy to reduce *Streptococcus mutans* in dental caries.¹⁷ They concluded that this treatment can be used as a photosensitising agent for photodynamic therapy to reduce the *S. mutans* burden in dental caries.

Laser dentistry offers many application options and numerous research approaches that might be interesting to investigate or to stay up-to-date with for practitioners. This consideration of recent literature has shown that there is still much potential for the increased use and application of lasers in the different fields of dentistry.



contact

Dr Igor Cernavin

Prosthodontist
 Honorary Senior Fellow at the
 University of Melbourne School of Medicine,
 Dentistry and Health Sciences
 Director and co-founder of the
 Asia Pacific Institute of Dental Education
 and Research (AIDER)
 Australian representative of WFLD

Private practice
 274 Main Rd East
 St Albans VIC 3021, Australia



Kurz & bündig

Zahnmedizinern, welche Laser zukünftig in ihrer Praxis einsetzen möchten oder wissenschaftliche Erkenntnisse zu den passenden Einsatzgebieten suchen, bieten aktuelle Studien viele Einblicke in nutzbringende Anwendungsbereiche. Der Autor stellt diverse aktuelle Studien der Laserforschung und -nutzung wie zum Beispiel zur Verwendung von Lasern zur Implantatfreilegung oder zur Auswahl des geeigneten Lasers für das persönliche Tätigkeitsfeld vor. Kurze Auszüge und Zusammenfassungen geben Anwendern einen ersten Eindruck und bieten ihnen eine Übersicht zu den gegenwärtigen Einsatzmöglichkeiten und Forschungsansätzen, u. a. zum Einsatz der Lasertherapie zur Minimierung oraler Beschwerden von Krebspatienten. Der Autor offeriert dem Leser verschiedene Optionen für die interessante Lektüre oder weiterführende Recherche.

Die Laserzahnmedizin bietet viele Anwendungsoptionen, sowie zahlreiche interessante Forschungsgebiete, bei welchen es sich für Anwender lohnt, auf dem neusten Stand zu bleiben. Die hier aufgezeigte Literatur verdeutlicht, dass noch sehr viel Potenzial zum vermehrten Einsatz von Lasertechnik in den diversen Feldern der Zahnmedizin besteht.




LIGHT INSTRUMENTS
RISE ABOVE TECHNOLOGY

NEW!

New Endo Tip for LiteTouch™

Get the best results with the most efficient technology for cleaning and disinfecting the root canal system.

- Introducing the brand new **Endo Tip** for LiteTouch™, especially suitable for cleaning and disinfecting the root canal system during endodontic treatment and retreatment.
- The new **Endo Tip** is a flexible, hollow, side-firing conical tip, with circumferential spiral slits located throughout the length of the tip. The tip is designed to fit the shape and volume of root canals prepared by NiTi rotary instrumentation. It emits the Er:YAG laser radiation laterally, throughout all the root canal walls.
- The tip is sealed at its end, preventing the transmission of radiation through the apical foramen of the tooth.
- The new **Endo Tip's** special design allows omnidirectional (360°) laser beam delivery throughout the tip's length, rather than direct emission through a single opening at the tip end. This unique design enables more efficient cleaning, disinfection and removal of smear layer.



SEM studies of extracted teeth after using the new **Endo Tip** showed clean surfaces of the root canal walls, free of smear layer and debris.

The **Endo Tip** was developed by Prof. Adam Stabholz and Light Instruments. The SEM studies were conducted at the Hebrew University - Hadassah School of Dental Medicine in Jerusalem, by Prof. Adam Stabholz & Dr. Sharonit Sahar-Helft.

LiteTouch™, the world's most versatile non-fiber Er:YAG laser device for soft and hard tissue dental treatments.

To learn more about our innovative technologies please visit:

www.light-inst.com

