

# implants

international magazine of oral implantology



## research

The key role of vitamin D

## case report

Immediate rehabilitation

## industry

Aesthetics for all intends and purposes





# Guided Tissue & Bone Regeneration



## CERASORB® M

Resorbable, pure-phase  
 $\beta$ -tricalcium phosphate



## CERASORB® Paste

Resorbable  $\beta$ -tricalcium  
phosphate paste +  
hyaluronic acid matrix



## CERASORB® Foam

Resorbable  $\beta$ -tricalcium  
phosphate + collagen  
matrix



## Inion®

Resorbable pin  
fixation system



## Osgide®

Resorbable  
polylactic acid (PLA)  
membrane



## Epi-Guide®

Resorbable  
polylactic acid (PLA)  
membrane



## Ti-System

System for  
fixation of foils  
and membranes



## Tefguide®

Non-resorbable  
PTFE membrane



## Osbone®

Synthetic  
spongy bone  
substitute



## Stypro®

Resorbable  
hemostyptic

# curasan

Regenerative Medicine

Phone: +49 6027 / 40 900-0

[info@curasan.com](mailto:info@curasan.com)

[www.curasan.com](http://www.curasan.com)

# One toolbox for your needs!



Dr Rolf Vollmer

First Vice President and Treasurer of DGZI



## The right measures at **the right time**

It was certainly one of the more memorable press conferences in recent dental history that Henry Schein held online on late Thursday afternoon at the end of February. Stanley M. Bergman, chairman of the board and CEO of Henry Schein, elaborated on questions about moral duties and entrepreneurial responsibility in times of a global health crisis. All over the world, people are losing trust in their governments, according to Bergman. It has become apparent that businesses are able to react more swiftly to the challenges posed by the pandemic than are governmental institutions owing to their innovation strength and flexibility. Procurement of validated personal protective equipment and especially vaccine development come to mind here. Bergman spoke in favour of global efforts to improve material and infrastructural requirements of healthcare systems to better fight the current and future pandemics.

In Germany, political entities and expert dental associations quickly switched to crisis mode, implementing improved hygiene guidelines in order to maintain dental care during lockdown. In an unprecedented response, it was the dental practices themselves who streamlined their patient management, readjusted their already outstanding hygiene protocols and implemented infection protection in shift operations, and thus strongly consolidated patients' trust in dental care. In doing so, Germany-based dentists set an international example.



The latest numbers from the German statutory occupational accident insurance association for non-state healthcare and welfare institutions in the Berufsgenossenschaft für Gesundheitsdienst und Wohlfahrtspflege (BGW) prove that, not only patients, but also practice staff are well protected against infections by consistent hygiene measures. In 2020, the BGW recorded 19,774 reportable occupational suspected cases of SARS-CoV-2—only 85 of which were related to the dental profession. Of course, these figures only hold true for Germany. Nevertheless, they bolster dental offices worldwide, because they indicate that, for dentists and their staff, a sustainable way out of the crisis with comparatively few infections is possible by appropriate crisis management in combination with a high standard of hygiene.

In this spirit, I wish you pleasant reading of the new **implants—international magazine of** oral implantology, continuously safe practice management and a great deal of courage to support your patients. After all, implantology is an integral part of sustainable healthcare!

Yours,

A handwritten signature in black ink, appearing to read 'Dr Rolf Vollmer'.

Dr Rolf Vollmer



Cover image courtesy of Fotona d.o.o.  
www.fotona.com



## editorial

- The right measures at **the right time** 03  
Dr Rolf Vollmer

## research

- The key role of **vitamin D** in immune health and regeneration 06  
Prof. Shahram Ghanaati, Dr Karl Ulrich Volz & Dr Sarah Al-Maawi

## case report

- Immediate rehabilitation of a **completely edentulous** maxilla 12  
Dr Jean-Baptiste Verdino
- Effectively **designing** the aesthetic zone 16  
Prof. Bilal Al-Nawas, Dr Keyvan Sagheb,  
Dr Stefan Wentaschek & Dr Yasmin Habibi
- Pre-prosthetic periodontal plasty with the **Er:YAG** laser 20  
Dr Fabrice Baudot

## industry

- Aesthetics** for all intents and purposes 24  
Dipl.-Ing. Axel Reichert
- Making your work **flow** in implant dentistry 26  
Dentsply Sirona, Sweden
- Low CHX concentrations with additives **effective against** biofilm 28  
Curaden, Switzerland
- Implant integration made **predictable** 30  
Prof. David L. Hoexter

## news

- manufacturer news 34
- news 40

## about the publisher

- imprint 42

# FLEX YOUR MUSCLES

OUR SHORTCUT TO IMMEDIACY:  
CONELOG® PROGRESSIVE-LINE FLEX



## PROSTHETIC SIMPLICITY AND CLEVER TRAY

- Consequently prosthetically oriented implant system
- Small tray with all instruments you need for implant placement
- Washtray function: validated for reprocessing
- Implant design for high stability in soft bone

## EXPERIENCE THE PROGRESSIVE EFFECT.



video animation

CURIOUS? PHONE +49 7044 9445 239

[sales@camlog.com](mailto:sales@camlog.com)

[www.camlog.com](http://www.camlog.com)

**camlog**

 **biohorizons  
camlog**


# The key role of vitamin D in immune health and regeneration

## The evidence for supplementation

Prof. Shahram Ghanaati, Dr Karl Ulrich Volz & Dr Sarah Al-Maawi, Germany & Switzerland

**A healthy immune system** is the basis of general good health and a good immune defence. It has been proved that individual habits, nutrition and the environment have an influence on our health.<sup>1</sup> A balanced and healthy diet in particular is the key to a healthy human body. An unbalanced diet can seriously impair the immune system and increase the risk of chronic disease as a result.<sup>1</sup> In the last decade, chronic diseases such as diabetes mellitus,

obesity and cardiovascular disease have surged sharply in various countries. A major reason for this is an increasingly unhealthy living environment and increasingly unhealthy lifestyle choices, especially in industrialised countries.<sup>2</sup> The role of food components and especially vitamins has become increasingly important in various areas. In 1928, the German biochemist Adolf Windaus was awarded the Nobel Prize in Chemistry for his work on the correlation between sterols and vitamins, which sparked further research interest in vitamin D.<sup>3</sup>



Vitamin D can be produced in a physiological way in the human body. Sunlight is essential for this endogenous synthesis, which takes place primarily in the skin, where 7-dehydrocholesterol is converted into cholecalciferol (vitamin D<sub>3</sub>) by UVB rays. In order to reach its biologically active form, cholecalciferol undergoes further conversion steps in the liver (calcidiol) and in the kidney (calcitriol). The latter is the biologically active form of vitamin D and acts as a transcription factor. After binding to the vitamin D receptor, calcitriol regulates the expression of various proteins in the cell. The physiological mode of action of calcitriol therefore resembles that of a hormone and not that of a vitamin. That is why vitamin D, as a precursor of calcitriol, should rather be regarded as a prohormone (Fig. 1).<sup>4,5</sup> The connection between vitamin D and parathyroid hormone was recognised shortly after its discovery. Within this context, the regulatory effect of vitamin D on the mineral balance of the body and in particular the regulation of calcium and phosphate levels was emphasised.<sup>6-8</sup> Furthermore, it was established quite early on that vitamin D plays an important role in mineralisation and bone formation. Consequently, many studies have focused on the influence of vitamin D on skeletal health and the treatment of diseases such as osteoporosis. These findings have contributed to vitamin D being primarily associated with bone health in the public perception.

However, some studies have shown the positive effect of vitamin D on the immune system too and thus on the general health of the body. Several studies have shown that vitamin D has a preventive effect on chronic diseases such as diabetes mellitus, hypertension and cardiovascular dis-

ease.<sup>9</sup> Studies also report its potential anti-inflammatory and antiviral effects.<sup>10</sup> In this context, it has been shown that vitamin D supplementation in pupils could reduce the incidence rate of influenza virus infection.<sup>11</sup> These rather new findings and the immunomodulatory effects of vitamin D demonstrate the importance of maintaining healthy vitamin D levels in the body. Since endogenous vitamin D synthesis is compromised by relatively short exposure to sunlight in most countries, the need for exogenous supply is becoming increasingly important. However, the intake of vitamin D through food seems to be insufficient in the general population, which has contributed to a global vitamin D deficiency pandemic.<sup>12</sup> This pandemic has already been documented and reported in numerous studies in various countries.<sup>13</sup> Nevertheless, its importance is still mostly under-estimated in most countries.

The concept of supplementation with vitamin D preparations was first introduced in the 1940s. Today, 90 years later, there are still no uniform recommendations regarding the dose to be taken. One of the reasons for this is the historical development and the association of vitamin D with bone health and the new knowledge about its further extensive capabilities. Although there is a growing amount of data on the non-skeletal effects of vitamin D and its preventive role in many chronic diseases, current dose recommendations are still based solely on bone re-

quirements. Another issue is the difficulty in standardising methods for the determination of serum vitamin D levels. This review therefore focuses on the non-skeletal effects of vitamin D and its supplementation dose based on randomised controlled clinical trials. It provides an overview of the new findings and treatment protocols.

## Immune system booster in the case of chronic and infectious disease

There is increasing interest in the study of the immune system-supporting mechanisms of vitamin D. Interestingly, the majority of body cells express vitamin D receptors on their surfaces, which emphasises the multimodal action of vitamin D. Owing to its regulatory effect, the active form of vitamin D as a hormone can intervene in the synthesis of various cytokines and regulate them according to their condition.<sup>14</sup> It has been shown that vitamin D inhibits the production of pro-inflammatory cytokines, whereas it up-regulates the synthesis of anti-inflammatory signal molecules.<sup>5</sup> In this way, it exerts its immunomodulatory effect and supports the differentiation of lymphocytes into Th2 cells and regulatory T cells.<sup>14</sup> This could explain its potential preventive influence in chronic and infectious diseases. However, these mechanisms of action still remain largely unexplained for the respective indications.

AD

## SUBSCRIBE NOW!

implants – international magazine  
of oral implantology

Interdisciplinary, transparent & market orientated



ONLINE  
SUBSCRIPTION HERE



www.oemus-shop.de

Fax: **+49 341 48474-290**

I would like to subscribe for the following journals:

- ☐ ceramic implants — international magazine of  
ceramic implant technology 2 issues p.a. € 30\*
- ☐ implants — international magazine of  
oral implantology 4 issues p.a. € 44\*

Terms & Conditions: The subscription may be cancelled in written form without due justification within 14 days of order by contacting OEMUS MEDIA AG, Holbeinstraße 29, 04229 Leipzig, Germany. Dispatching notification in good time will suffice. The subscription is automatically extended by another twelve months if it is not cancelled in written form 6 weeks prior to the end of the reference period.

\* All prices include VAT, shipping and handling.

Last Name, First Name

Company

Street, ZIP, City, Country

E-mail

Credit Card Number

Expiration Date

Security Code

Stamp

Signature

**OEMUS MEDIA AG**

Holbeinstraße 29 · 04229 Leipzig · Germany · Phone: +49 341 48474-201 · s.schmehl@oemus-media.de



The correlation between vitamin D levels and the prevalence of various chronic diseases has been shown in several clinical studies. A meta-analysis of 25 prospective cohort studies has shown that low vitamin D levels increase the risk of developing cardiovascular disease. In about 10,000 patients, the risk of cardiovascular disease was about 44 % higher than in people with healthy vitamin D levels.<sup>15</sup> Another study showed a correlation between vitamin D levels and the development of hypertension. It examined 8,155 patients suffering from hypertension and vitamin D deficiency. After the vitamin D deficiency had been eliminated, 71 % of the patients no longer showed any symptoms or had measurably high blood pressure.<sup>16</sup> A positive influence of vitamin D has also been demonstrated in the development of Type 2 diabetes mellitus. It was shown that the number of patients in a prediabetic stage and with a vitamin D deficiency was significantly lower than in the untreated group, once the vitamin D deficiency had been eliminated.<sup>17</sup>

Furthermore, the potential of an anti-infectious or antiviral effect of vitamin D has been increasingly investigated in recent years. As a result, vitamin D has gained greater significance as a preventive or adjuvant therapy.<sup>11,18</sup> A systematic review has shown that a vitamin D deficiency is associated with a higher viral load in hepatitis B patients.<sup>19</sup> Furthermore, it was shown that vitamin D can inhibit a herpesvirus infection through its anti-inflammatory and supportive defence effect.<sup>20</sup> In addition, studies have shown that vitamin D supplementation reduces the prevalence of influenza infections during influenza outbreaks.<sup>21</sup> Another meta-analysis showed that the number of certain vitamin D receptor polymorphisms involved in processing of vitamin D correlates with an increased risk of a viral infection. Based on the vitamin D-mediated improved immune defence and its potential role as an antiviral agent,


its importance in the prevention of viral diseases is increasingly being investigated. Especially in the COVID-19 pandemic, vitamin D supplementation can play an important role in preventing and defeating infection.<sup>22</sup>

## Determination of vitamin D levels and definition of hypovitaminosis

Vitamin D is a lipophilic molecule that is transported in the blood by carrier proteins. Approximately 80 % of these molecules are bound to the vitamin D binding protein in this manner. A further 10–15 % are bound to albumin and the rest circulates freely in the blood. The determination of the vitamin D level as part of a routine examination involves measuring the total concentration of all these forms. The 25(OH)D serum concentration is widely recognised as a reliable marker of vitamin D levels.<sup>12</sup> Similar to other vitamins and blood components, the vitamin D concentration is usually expressed in nanograms per millilitre (ng/ml) or in nanomoles per litre (nmol/l). Both units are used, depending on the individual testing laboratory. Here, it must be noted that 1 nmol/l equals 0.4 ng/ml. The definition of a healthy vitamin D level and thus hypovitaminosis is a matter of much debate. In the literature, a vitamin D level of less than 30 ng/ml (75 nmol/l) is considered a vitamin D deficiency (hypovitaminosis).<sup>13,19,23,24</sup> In various countries, studies have reported a general vitamin D deficiency. Observational studies have documented that the prevalence of vitamin D levels of below 20 ng/ml (50 nmol/l) is as much as 24 % in the US, 37 % in Canada and 40 % in Europe.<sup>13,24</sup> The German Robert Koch Institute reported that 58 % of 18- to 79-year-olds in Germany have a level of below 20 ng/ml (50 nmol/l).<sup>25</sup> This vitamin D deficiency pandemic was recognised as such several years ago. However, not much has been done in terms of supplementation and defining a sufficient dose. A pilot study examined the vitamin D levels of medical staff in the clinic for oral and maxillofacial plastic surgery at Goethe University in Frankfurt am Main in Germany. Out of 24 participants, 85.7 % had a vitamin D deficiency with a value below 30 ng/ml, whereas 45.8 % even had a value of below 10 ng/ml (Fig. 2). It is important to emphasise that a healthy vitamin D value is considered to be between 40 ng/ml and 60 ng/ml.

## Current guidelines for vitamin D supplementation

Given that, in most cases, endogenous synthesis of vitamin D is insufficient owing to limited exposure to sunlight, the body's vitamin D intake should also come from food or dietary supplements. The amount of vitamin D absorbed can be expressed in two units: micrograms (µg) and international units (IU). One microgram equals 40 international units (1 µg equals 40 IU). These units



**Fig. 2:** Distribution of vitamin D levels according to a pilot study conducted by the Clinic for Oral and Maxillofacial Plastic Surgery at Goethe University Frankfurt am Main.



must be considered when administering vitamin D. Since in most cases vitamin D intake via food is insufficient for the body's needs, supplementation with vitamin D preparations is an utmost necessity. In the literature, the current recommendations for doses to be administered are largely inconsistent and are mainly based on the estimated requirements of maintaining optimal bone health. The recommendations range from 400 IU/day to 4,000 IU/day. The European Food Safety Authority recommends

a dose of 600 IU/day for healthy adults.<sup>22</sup> A similar recommendation, a dose of 400 IU/day, has been published by the Scientific Advisory Committee on Nutrition in the UK.<sup>26</sup> The Institute of Medicine Committee in the US recommends a dose of 600 IU/day for adults under 70 years of age and a dose of 800 IU/day for those over that age.<sup>27</sup> The American Association of Clinical Endocrinology recommends a dose of 1,000–4,000 IU/day.<sup>28</sup> The recently updated reference values of 2012 from the German

| Category                                                                                   | Dose                            | Administration duration | Initial concentration | Targeted concentration | Side effects                                                              |
|--------------------------------------------------------------------------------------------|---------------------------------|-------------------------|-----------------------|------------------------|---------------------------------------------------------------------------|
| Prevention in pupils <sup>21</sup>                                                         | 1,200 IU/day                    | 12 months               | Not specified         | Not specified          | None                                                                      |
| Cancer, cardiovascular disease <sup>30</sup>                                               | 2,000 IU/day                    | 12 months               | 29.8 ng/ml            | 41.8 ng/ml             | None                                                                      |
| Diabetes mellitus <sup>17</sup>                                                            | 4,000 IU/day                    | 12 months               | 28.0 ng/ml            | 52.3 ng/ml             | None                                                                      |
|                                                                                            | 4,000 IU/day                    | 24 months               | 28.0 ng/ml            | 54.3 ng/ml             | None                                                                      |
| Ventilated patients in intensive care <sup>31</sup>                                        | 50,000 IU/day                   | 5 days                  | 23.2 ng/ml            | 45.0 ± 20.0 ng/ml      | None                                                                      |
|                                                                                            | 100,000 IU/day                  | 5 days                  | 20.0 ng/ml            | 55.0 ± 14.0 ng/ml      | None                                                                      |
| Test persons with a vitamin D deficiency <sup>32</sup>                                     | 25,000 IU/fortnight             | 2 months                | 7.6 ng/ml             | 19.0 ng/ml             | None                                                                      |
|                                                                                            | 25,000 IU/week                  | 1.5 months              | 8.0 ng/ml             | 25.0 ng/ml             | None                                                                      |
|                                                                                            | 25,000 IU/week                  | 2 months                | 8.4 ng/ml             | 35.6 ng/ml             | None                                                                      |
| Test persons with a vitamin D deficiency <sup>33</sup>                                     | 1,000 IU/day                    | 5 months                | 28.8 ng/ml            | 33.6 ng/ml             | None                                                                      |
|                                                                                            | 5,000 IU/day                    |                         | 27.0 ng/ml            | 64.0 ng/ml             | None                                                                      |
|                                                                                            | 10,000 IU/day                   |                         | 26.0 ng/ml            | 89.6 ng/ml             | None                                                                      |
| Breast cancer patients with bone metastasis <sup>34</sup>                                  | 7,000 IU/day                    | 4 months                | < 20.0 ng/ml          | Not specified          | None                                                                      |
| Psychiatric clinic <sup>24,35</sup>                                                        | 5,000 IU/day                    | 12 months               | 24.0 ng/ml            | 68.0 ng/ml             | None                                                                      |
|                                                                                            | 10,000 IU/day                   | 12 months               | 25.0 ng/ml            | 96.0 ng/ml             | None                                                                      |
| Test persons with a vitamin D deficiency <sup>36</sup>                                     | 100,000 IU/month (3,000 IU/day) | 36 months               | 24.4 ng/ml            | 54.0 ng/ml             | None                                                                      |
| Multiple sclerosis <sup>37</sup>                                                           | 20,000 IU/day                   | 12 months               | 21.6 ng/ml            | 44.0 ng/ml             | None                                                                      |
| Multiple sclerosis <sup>38</sup>                                                           | 50,000 IU/week (7,142 IU/day)   | 6 months                | 15.3 ng/ml            | 33.7 ng/ml             | None                                                                      |
| Asthma, rheumatic arthritis, rickets, tuberculosis in the 1930s and 1940s <sup>24,39</sup> | 60,000–600,000 IU/day           | Not specified           | Not specified         | Not specified          | Hypercalcaemia as a result of over-physiological vitamin D concentrations |

**Table 1:** Overview of the vitamin D doses administered in selected randomised clinical studies.