

roots

the international C.E. magazine of endodontics

4²⁰¹⁴

_C.E. article


From everyday dentistry to advanced photoacoustic endodontic applications (PIPS)

_education

Control the anatomy; control procedural training

_association news

Peter S. Weber is named executive director of AAE



FOCUS ON PRECISION

With over 69 years of optical experience, Seiler Instrument is a worldwide leader in the field of Microscopy.



**FREE
30 DAY
DEMO**

Introducing the all New Evolution XR6 LED: With over 80,000LUX, 5800K for a pure daylight look and over 50,000 hours of bulb life, you will see clearer, sharper images every time.

In addition to superior clarity, the XR6 also offers all APOchromatic Lenses, a new ergonomic design for an almost effortless movement, and comes at a highly competitive price.

Don't be left in the dark, try a Seiler Microscope today!

Come visit Seiler at the GNYM and test drive the NEW LED Microscope at BOOTH #5200.



**SEILER**
PRECISION MICROSCOPES
Providing High Quality Optics since 1945
800-489-2282 www.seilermicro.com

Exploring a world of knowledge



Fred Weinstein, DMD, MRCD(C),
FICD, FACD

There is always plenty to learn in our specialty.

In this issue of *roots*, you can find many helpful articles, including a report on the use of dual wavelength lasers by Dr. Lawrence Kotlow, Dr. Enrico DiVito and Dr. Giovanni Olivi. As their article points out, these lasers have uses in many different clinical situations. Perhaps most impressive is the use of this equipment in root canal therapy. The images on the cover of this issue attest to the effectiveness of lasers in cleansing the anatomy.

Also in this issue of *roots*, you can learn about the 3-D training replicas available from Dr. L. Stephen Buchanan and his team at Dental Education Laboratories. These tooth models offer many advantages for those who wish to hone their skills.

Every issue of *roots* magazine also contains a C.E. component. By reading the article on dual wavelength lasers mentioned above, and then taking a short online quiz about this topic at www.DTStudyClub.com, you will gain one ADA CERP-certified C.E. credit.

To learn more about how you can take advantage of this and many other C.E. opportunities, visit www.DTStudyClub.com. You need only register at the Dental Tribune Study Club website to access these C.E. materials free of charge. You may take the C.E. quiz after registering on the DT Study Club website.

I know that taking time away from your practice to pursue C.E. credits is costly in terms of lost revenue and time, and that is another reason *roots* is such a valuable publication. I hope you will enjoy this issue and that you will take advantage of the C.E. opportunity.


Finally, I would like to mention that with this publication, we are merging the North American and international editions of *roots*. Welcome to all readers, wherever you are from.

As always, I welcome your comments and feedback.

Sincerely,

A handwritten signature in cursive script that reads "Fred".

Fred Weinstein, DMD, MRCD(C), FICD, FACD
Editor in Chief



| C.E. article

- 06 From everyday dentistry to advanced **photoacoustic endodontic applications (PIPS)**: Er:YAG & Nd:YAG dual wavelength laser
_Lawrence Kotlow, DDS, Enrico DiVito, DDS & Giovanni Olivi, MD, DDS

- 22 An industry-first solution from Planmeca: **Imaging, CAD/CAM and dental units** in one software
- 24 FKG revolutionizes endodontics with the **BT-Race sequence** for a biological and conservative treatment

| education

- 12 Control the anatomy; control **procedural training**
_L Stephen Buchanan, DDS, FICD, FACD

| about the publisher

- 26 _imprint

| industry education

- 16 **LVI Core I** three-day course is designed for doctors and their teams to learn together
_Mark Duncan, DDS, FAGD, LVIF, DICOI, FICCMO

| associaton news

- 17 Peter S. Weber is named executive director of **AAE**




| industry

- 18 **How much money** are you wasting?
- 20 **GentleWave**: Multisonic Ultracleaning System

| on the cover

Root canal dentinal walls after PIPS laser treatment, shown under magnification. See article beginning on page 6.



TrueTooth®

Not models. Replicas of actual teeth.

"The real value in the TrueTooth Replicas is that our students can appreciate the complexity of the root canal system. We have the benefits of students working on replica teeth that we select, and everyone is working on the same tooth. It levels the playing field and takes away the infection control concerns. We can concentrate our energy on technique instead of tooth selection."

Donald J. Kleier, D.M.D.
Professor and Chair,
Division of Endodontics, School of Dental Medicine, University of Colorado



Replica #3-001

*Now available
in accessed
and shaped
versions!*



Replica #9-001

"No more explaining to students how the anatomy is different than natural teeth. Get real, people."

Ronald R. Lemon, D.M.D.
Coordinator of Predoctoral Endodontic Program and Associate Dean for Advanced Education
School of Dental Medicine, University of Nevada, Las Vegas

"TrueTooth Replicas brought anatomic realism to our preclinical endodontics course!"

Division of Endodontics,
University of Minnesota, School of Dentistry

Almost as Good as the Real Thing

Available exclusively through Dental Education Laboratories
Your Premier Resource for Endodontic Continuing Education

800.528.1590 / 805.899.4529 / www.DELendo.com Patent Pending

From everyday dentistry to advanced photoacoustic endodontic applications (PIPS): Er:YAG & Nd:YAG dual wavelength laser

Authors_Lawrence Kotlow, DDS, Enrico DiVito, DDS, and Giovanni Olivi, MD, DDS

_c.e. credit

This article qualifies for C.E. credit. To take the C.E. quiz, log on to www.dtstudyclub.com. Click on 'C.E. articles' and search for this edition (Roots C.E. Magazine — 4/2014). If you are not registered with the site, you will be asked to do so before taking the quiz. You may also access the quiz by using the QR code below.



_Introduction

Lasers provide an exciting new technology that allows the dentist the ability to give patients optimal care without many of the "fear factors" found in conventional dental techniques. Used with proper understanding of laser physics, lasers are extremely safe and effective.

Using lasers for caries removal, perio treatment, endodontic treatment, bone management, cutting and shaping, and soft-tissue procedures can reduce postoperative discomfort and infection, and provide safe, simple in-office treatment. As a result, we can improve our efficiency, expand what we can do, achieve better results and increase production.



Lasers represent a real quantum leap forward in the treatment of our patients, including the pediatric

patient. The U.S. Food and Drug Administration (FDA) gave approval for the use of the Er: YAG laser in 1997 for both hard- and soft-tissue procedures. The erbium doped (erbium particles placed within the YAG crystal) crystal of Yttrium-Aluminum-Garnet's (Er:YAG) development and success has made the treatment of children safer and quicker.

Plainly stated, a laser is a piece of equipment that creates a concentrated monochromatic beam of visible or infrared light that can be absorbed by a specific target. Since then, laser-assisted dental care has changed forever the way dentists can prepare diseased teeth, ablate bone and treat soft-tissue abnormalities and disease. An entire new standard of care is becoming a reality.

Lasers and pediatric dentistry are a perfect fit. There are a wide range of hard and soft dental pro-

Figs. 1, 2 Representative sample images of root canal dentinal walls irrigated with 17 percent EDTA and PIPS for 20 seconds. (Photos/ Provided by Technology4Medicine)



cedures that may be completed using lasers as an alternative to conventional dental care on adults and, especially, children. Many of these procedures may be treatments dentists historically refer out to other specialists; however, if you understand and use your laser efficiently, you will discover that many of these are procedures that every dentist can easily complete.

The question that is often the major concern and barrier to investing in lasers is the how this investment will pay for itself, more recently described as return on investment (ROI). Will it pay for itself? We prefer to speak of this as the secondary effect. If you understand your laser, it will easily pay premiums on your investment, and the cost factor becomes a non-issue.

The purchasing of lasers is an investment, not an expense, for any dental practice.

Lasers represent a fundamental change in the entire way dentistry has been taught. We can now rethink and often modify G.V. Black's principle of extension for prevention with the concept of minimally invasive micro-dentistry. We need to understand that laser dentistry is one portion of an entire new way of practicing conservative, pain-free dentistry.

The laser that we call the "all-purpose" laser is the Lightwalker Er:YAG & Nd:YAG laser, manufactured by Fotona and distributed in the United States by Technology4Medicine. The Er:YAG produces its effect at 2940 nm and has as its primary tissue target water and hydroxyapatite. It is very safe, relatively quiet, eliminates the smells and vibrations associated with the dental handpiece and, most importantly, is much more comfortable for the patient, significantly reducing the need for local anesthesia.

The use of the new generation erbium lasers for repair of incipient hard-tissue disease allows the dentist to provide a stress-free means of restoring teeth in a minimally invasive manner, most often with no shot and no numb lip, without the need for any local anesthetics.

The erbium laser can be used for restoring primary and permanent teeth, eliminating or reducing the amount of local anesthetics. In most cases, the patient will not require numbing for Class 1, 2 (sometimes), 3, 4, 5, 6 restorative procedures using bonded restorative materials. Using the concept of minimally invasive restorative procedures, the Er:YAG laser allows the operator to remove only diseased tissue and thus preserves much more of the healthy, unaffected tooth.

In cases where alloy is preferred, the laser's analgesia effect may also allow the dentist to create a restorative preparation using a conventional handpiece that is not meant for bonding. The erbium laser is effective because of its effect on its target, water within the tooth structure. This effect occurs when the laser heats up water within the target

tissue, causing it to create small microscopic explosions (photothermal followed by photoacoustical effects). When applied to soft tissue, bone or teeth and cavities, the explosions then cause the areas to be vaporized.

Er:YAG laser 2940 nm: Soft-tissue procedures

There is a wide array of soft-tissue procedures that can be completed using the all-purpose laser: maxillary and mandibular frenum revisions, lingual frenum revisions, treatment of pericoronal pain or infection, removal of hyperplastic tissue because of drugs or poor oral care in orthodontic patients, biopsies, treatment of aphthous ulcers and herpes labialis, pulpotomies, removal of impacted teeth and, in adults, apicoectomies and bone recontouring.

Pulpotomies

Parents often express concern about the need to take radiographs because of the nature of X-rays and their possible side effects on a child's overall health. They question the use of alloys because of the chemical makeup of the alloy. Whether these should be a real concern in today's dental care is open to debate, depending on your individual beliefs. There are also concerns by many, although not as loudly, about the effect of various pulpotomy procedure medicaments used in pulpotomy procedures, such as formocresol.

Lasers provide a safe, non-chemical, effective and alternative treatment for pulpotomies. During the span of eight years, post-treatment results on more than 4,000 pulpotomies using the erbium (2940 nm) laser provide ample evidence that this method is both effective and safe for children without the need for introducing chemicals or using electro-surgery methods.

When the final result of orthodontic positioning of the front teeth results in gingival hypertrophy, the laser can be a useful tool to increase crown length and give the patient a more esthetic smile. This may often be accomplished without the need for local anesthesia. Patients who have medically induced hyperplastic tissue, such as patients requiring dilantin, can also have their tissue reduced and reshaped with the erbium.

In addition to the many examples described in this article, lasers can be used for additional procedures not usually required in pediatric dentistry, such as revisions of the abnormal mandibular frenum, often avoiding the need for soft-tissue grafts, crown-lengthening procedures where bone requires recontouring, apicoectomies, removal of bony exostoses, removal of third molar impactions, removal of root remnants, incising and draining soft-tissue

infections, advanced periodontal treatments and the latest in advanced endodontic treatment via photon-induced photoacoustic streaming.

Photoacoustic endodontics using PIPS

The goal of endodontic treatment is to obtain effective cleaning and decontamination of the smear layer, bacteria and their byproducts in the root canal system. Clinically, traditional endodontic techniques use mechanical instruments, as well as ultrasonic and chemical irrigation, in an attempt to shape, clean and completely decontaminate the endodontic system but still fall short of successfully removing all of the infective microorganisms and debris. This is because of the complex root canal anatomy and the inability for common irrigants to penetrate into the lateral canals and the apical ramifications. It seems, therefore, appropriate to search for new materials, techniques and technologies that can improve the cleaning and the decontamination of these anatomical areas.

Among the new technologies, the laser has been studied in endodontics since the early 1970s¹⁻³ and has become more widely used since the '90s.⁴⁻⁶

Different wavelengths have been shown to be effective in significantly reducing the bacteria in the infected canals, and important studies have confirmed these results *in vitro*.⁷ Studies reported that near infrared laser are highly efficient in disinfecting the root canal surfaces and the dentinal walls (up to 750 microns for the diode 810 nm and up to 1 mm for the Nd:YAG 1064 nm). On the other hand, these wavelengths did not show effective results in debriding and cleansing the root canal surfaces and caused characteristic morphological alterations of the dentinal wall. The smear layer was only partially removed and the dentinal tubules primarily closed as a result of melting of the inorganic dentinal structures.^{5,8}

Other studies reported the ability of the medium infrared laser in debriding and cleaning root canal walls.^{9,10} The bacterial load reduction after erbium laser irradiation demonstrated high on the dentin surfaces but low in depth of penetration because of the high absorption of laser energy on the dentin surface.⁷ Also the laser activation of commonly used irrigants (LAI) resulted in statistically more effective removal of debris and smear layer in root canals compared with traditional techniques (CI) and ultrasound (PUI).^{11,12} Additionally, the laser activation method resulted in a strong modulation in reaction rate of NaOCl, significantly increasing production and consumption of available chlorine in comparison to ultrasound activation.¹³

A recent study has reported how the use of an Er:YAG laser, equipped with a newly designed radial and stripped tip, in combination with 17 percent EDTA solution, using very low pulse duration (50 mi-

croseconds) and low energy (20 mJ) resulted in effective debris and smear layer removal with minimal or no thermal damage to the organic dentinal structure through a photoacoustic technique called photon induced photoacoustic streaming or "PIPS."^{14,15} Also the same photoacoustic protocol in combination with 5.25 percent sodium hypochlorite solution has been investigated and shown to reduce the bacterial load and its associated biofilm in the root canal system three dimensionally.¹⁶

Other similar studies are in progress for publication and the results are promising and suggest a three-dimensional positive effect of this laser-activated decontamination (LAD) method.

The purpose of this article is to present briefly the experimental background of this laser technique and to introduce the clinical protocol.

Scientific background

The microphotographic recording of the LAI studies suggested that the erbium lasers used in irrigant-filled root canals generate a streaming of fluids at high speed through a cavitation effect.¹⁷ The laser thermal effect generates the expansion implosion of the water molecules of the irrigant solution, generating a secondary cavitation effect on the intracanal fluids. To accomplish this streaming, it is suggested the fiber be placed in the middle third of the canal, 5 mm from the apex and stationary.¹⁸ This concept greatly simplifies the laser technique, without the need to reach the apex and to negotiate radicular curves.

Also, the recorded video of the new technique, PIPS, showed a strong agitation of the liquids inside the canals. It differs from the already cited LAI technique by activating the irrigant solutions in the endodontic system through a profound photoacoustic and photomechanical phenomena. The use of low energy (50 microsecond pulse, 20 mJ at 15 Hz, 0.3 W average power, or less) generates only a minimal thermal effect. The study with thermocouples applied to the radicular apical third revealed only 1.2 degrees C of thermal rise after 20 seconds and 1.5 degrees C after 40 seconds of continuous radiation.¹⁴

When the erbium laser energy is delivered at only 50 microsecond pulse duration through a special designed tapered and stripped 400 microns tip (Fotona LightWalker, Technology4Medicine), it produces a large peak power of 400 watts when compared to a longer pulse duration. Each impulse, absorbed by the water molecules, creates a strong "shock wave" that leads to the formation of an effective streaming of fluids inside the canal while also limiting the undesirable thermal effects seen with other methodologies. The placement of the tip in the coronal portion only of the treated tooth allows for a more minimally enlarged




Fig. 3




Fig. 4

canal preparation with less thermal damage as seen with those techniques placed into the canal system.

The root canal surfaces irrigated with 17 percent EDTA and laser activated for 20 seconds showed exposed collagen matrix, opened tubules and the absence of smear layer and debris (Figs. 1-3). The rinsing with 5.25 percent sodium hypochlorite and laser irradiation for 20 seconds produced a strong activation of the solution, as reported by Macedo,¹³ improving the disinfecting action of the sodium hypochlorite.¹⁶ The disinfecting action of PIPS is very effective both on the root surface, the lateral canals and the dentinal tubules, as confirmed with SEM and confocal studies (Fig. 4).

The profound and distant effect of PIPS eliminates the need to introduce the tip into the root canal system. Unlike traditional laser techniques requiring placement of the tip 1 mm from the apex, or even 5 mm from the apex as proposed for LAI¹⁶, the PIPS tip is placed in the coronal portion of the pulpal chamber only and left stationary, allowing the photoacoustic effect to spread into the openings of each canal. A new tip design consisting of a 400-micron diameter, 12 mm long, tapered end is used for this technique (Fig. 5). The final 3 mm of coating is stripped from the end to allow for greater lateral emission of energy compared to the frontal tip.

This mode of energy emission allows for improved lateral diffusion with low energy and enhanced photoacoustic effect.

Discussion

Laser irradiation is a common technique used in endodontics to improve the cleaning, the debriding and disinfection of the root canal system. Many wavelengths and protocols are used. Near infrared lasers are used for the three-dimensional decontami-

nation of the endodontic system. Nd:YAG and diode lasers use thermal energy to destroy bacteria. Observations reveal a certain grade of thermal injury to the root canal surface and create a typical morphological damage. Moreover, they are not able to thoroughly remove the smear layer.

On the contrary, erbium lasers are used for their effective smear layer removal while their bactericidal activity is limited to the root surface. The placing of the tip close to the apex and its back movement during the activation process is related to the risk of apical perforation, ledging and surface thermal damage, because of the ablation ability of this wavelength. Also a combination of the near and medium infrared lasers has been proposed. A technique, called twilight endodontic treatment (TET), uses the erbium laser energy first, to clean the root canal surface and remove the smear layer, and the Neodimium:YAG laser second, used in dry mode as the final disinfecting step. All these techniques utilize traditional tips and fibers placed into the canal, close to the apex (1 mm) with all the corresponding thermal disadvantages observed in long, narrow and curve canals.

The erbium lasers are also used as a medium of activation of commonly used irrigants (LAI), avoiding the risk of thermal damage, while increasing the cleaning and disinfecting activity of the fluids. PIPS, in particular, reduces all these risks and disadvantages, thanks to the position of the tip in the coronal orifice only and to the use of minimally ablative energy levels of 20 mJ or less.

The findings of our studies demonstrated that PIPS technique resulted in a safe and effective debriding and decontaminating of the root canal system. Our clinical trials showed that PIPS technique greatly simplifies root canal therapy while facilitating the search for the apical terminus, debriding and maintaining patency.

Fig. 3 Representative sample image of root canal dentinal walls irrigated with 17 percent EDTA and PIPS for 20 seconds.

Fig. 4 SEM image of clean lateral canal.