

DENTAL TRIBUNE

— The World's Dental Newspaper • ASEAN Edition —



Eating disorders and dentistry: Identifying early signs in the dental office

Study analyses tooth survival after root canal therapy in US population

Dental sales a mixed bag as war and supply difficulties bog down pandemic recovery

In The Hot Seat

An Interview with Desmond Chong, the President of Malaysia Dental Industry Association (MDIA)

New Generation

Desktop 3D Printer for Digital Dentistry DentCare BQP-DI5

BenQ is the leader of DLP projector solution provider, that use this technology for the key part of DLP 3D printer at the same time. BQP-DI5 adopt DLP light curing technology of high resolution, that the advantage of high accuracy, large printing area, high working efficiency, and stability of working consumables.



SPEED or QUALITY? I WANT IT ALL!

- Made especially for dental application
- High speed, high accuracy, high stability and high durability
- Superior optical design to achieve excellent sharpness
- Z-axis liner motion guide provide high rigidity and stability
- The Z-axis creates high accuracy and high printing speed
- More user friendly and intuitive software

Unique design of resin tank

- Easy to detach the picker and tank
- Resin tank lifetime reaches 15 kg
(The lifetime depends on different resins)



BenQ BIS-1
Intraoral Scanner



Wireless Medical
Camera with Lamp



BenQ Lightweight
Reusable Instrument



AB Dental
Implant System



easyDU Preformed
Tooth Positioner

BenQ Integrated Digital Solutions

Solutions for implant, clear aligner and prosthetic treatment



BenQ
AB DentCare Corp



BenQ AB DentCare



Contact: ABD@BenQABDentcare.com / www.benqabdentcare.com

EDITOR'S MESSAGE

It's the month of July! We are grateful for your ongoing support for Dental Tribune ASEAN. As we are approaching the middle of the year, we've seen a few major events happening across the globe. On Jun 23, the World Health Organization declared the outbreak of monkeypox as a Public Health Emergency of International Concern due to the increasing number of cases which exceeds 17,000 in 75 countries.

Consumer Association of Penang (CAP) stated that stern action should be taken against fake dentists and inspections should be done by the authorities at dental clinics every six months. The unhygienic procedure performed by fake dentists is dangerous as many of them claim to have learned their skills from YouTube and such. It's also believed that they acquire the tools needed from hardware stores or suppliers from other countries that are deemed not intended for medical use.

On a side note, do check out our first-ever interview with Desmond Chong, the President of the Malaysia Dental Industry Association where we talked about how MDIA started its journey, milestones, and vision. MDIA also shared their outlook on Malaysia's




Susan Tricia
Editor

dental industry post-pandemic and matching grant worth RM10 million to develop the medical device industry to encourage investment and export market.

Dental Tribune ASEAN will be circulated in dental exhibitions, seminars, and even webinars across ASEAN providing additional mileage for advertisers of the magazine and reaching out to new subscribers. Visit our website at asean.dental-tribune.com for more updates. Let's do our part and get vaccinated.

Malaysia Publishing Partner

FBI Publications (M) Sdn Bhd 
Add: 9-3, Jalan PJU 5/6, Dataran Sunway, 47810, Petaling Jaya, Selangor
Tel: +603-6151 9178
Email: my@fireworksbi.com

International Headquarters

Publisher and Chief Executive Officer
Torsten R. Oemus

Chief Content Officer
Claudia Duschek

Dental Tribune International GmbH
Holbeinstr. 29, 04229 Leipzig, Germany
Tel.: +49 341 4847 4302 | **Fax:** +49 341 4847 4173

General requests
info@dental-tribune.com

Sales requests
ASEAN region:
vanny@fireworksbi.com
asean.dental-tribune.com

International:
mediasales@dental-tribune.com
www.dental-tribune.com

Publisher

FBI Publications (M) Sdn. Bhd.

Marketing Communication

Nur Izyan binti Dzulkifli
izyandzul@fireworksbi.com

Publication Manager

Vanny Lim
vanny@fireworksbi.com

Assistant Editor

Atthira Zawana
atthirah@fireworksbi.com

Creative Designer

Muhammad Fadzil
design@fireworksbi.com

Board of Directors

Kenny Yong
 Susan Tricia

Official Magazine of



Disclaimer

Material from Dental Tribune International GmbH that has been reprinted or translated and reprinted in this issue is copyrighted by Dental Tribune International GmbH. Such material must be published with the permission of Dental Tribune International GmbH. *Dental Tribune* is a trademark of Dental Tribune International GmbH. All rights reserved. © 2022 Dental Tribune International GmbH. Reproduction in any manner in any language, in whole or in part, without the prior written permission of Dental Tribune International GmbH is expressly prohibited. Dental Tribune International GmbH makes every effort to report clinical information and manufacturers' product news accurately but cannot assume responsibility for the validity of product claims or for typographical errors. The publisher also does not assume responsibility for product names, claims or statements made by advertisers. Opinions expressed by authors are their own and may not reflect those of Dental Tribune International GmbH.

Dental Tribune International GmbH, FBI Publications (M) Sdn Bhd nor its affiliated entities make any promises, implied or otherwise, about the fitness of any technique, piece of equipment, or material for a particular purpose.

Dental Tribune International GmbH, FBI Publications (M) Sdn Bhd, or its affiliated entities will not be liable for any damages of any kind or loss of profits that might arise from information found in this magazine regardless of whether Dental Tribune International GmbH has been advised of the possibility of such damages.

The material contained in this magazine is offered as information only and should not be used to replace any practice, financial, accounting, legal, or other professional advice that users might obtain from their own professional advisors.


Information in this magazine should not be substituted for the individual judgment brought to each clinical situation by a patient's dental practitioner. The techniques and opinions presented on this site reflect the science of dentistry at a particular point in time, and users should be aware that continued research on these topics may provide new knowledge or recommendations.

Dental Tribune International GmbH and FBI Publications (M) Sdn Bhd does not necessarily endorse the non-Dental Tribune International resources that may appear or be referenced in this magazine and offer no guarantee about the products or services they may provide.

Some sections in this magazine are published by separate organizations or entities, which are solely responsible for their content. Statements appearing in those portions are not necessarily endorsed by Dental Tribune International GmbH, FBI Publications (M) Sdn Bhd, or any of its subsidiaries. Also, Dental Tribune International GmbH and FBI Publications (M) Sdn Bhd does not offer any guarantee with respect to the accuracy of any items or claims contained therein.

Dental Tribune International GmbH and FBI Publications (M) Sdn Bhd does not, via this magazine or any of its affiliated media, encourage dentists to make any particular business decision based on issues addressed herein. Business decisions and decisions about individual patients and methods of practice are personal decisions, and the responsibility for these decisions rests upon the shoulders of the attending dental practitioner.

CONTENTS



Clinical news

6

Eating disorders and dentistry: Identifying early signs in the dental office

7

Study indicates denture wearers may be more at risk of nutritional deficiencies

World news

8

Study analyses tooth survival after root canal therapy in US population

10

Jönköping University aims to create new dental programme to address dentist shortage in Sweden

11

Parties to the Minamata Convention on Mercury further strengthen phase-down approach to dental amalgam

In the hot seat

12

An Interview with Desmond Chong, the President of Malaysia Dental Industry Association (MDIA)

Business

16

Dental sales a mixed bag as war and supply difficulties bog down pandemic recovery

18

Sales slowdown puts the brakes on Align Technology's record run

20

Digitally visualise clear aligner treatment results with new Align Technology outcome simulator

22

SmileDirectClub slashes net loss, signals major strategy change for 2022

24

Henry Schein to acquire Condor Dental and expand its operations in Switzerland





PURE SIMPLICITY



NEW CHIROPRO

IMPLANTOLOGY

NEW CHIROPRO PLUS

IMPLANTOLOGY

ORAL SURGERY

Control your entire implant and oral surgery motors using a single rotary knob. The new Chiropros from Bien-Air Dental have been designed around a single philosophy: simplicity.

Find out more at www.bienair.com

S W I S S  M A D E

Eating disorders and dentistry:

Identifying early signs in the dental office

By Iveta Ramonaite, Dental Tribune International



According to the Oral Health Foundation, dental teams can play a crucial role in identifying and acting on early signs of eating disorders. (Image: Paradise studio/Shutterstock)

According to some eating disorder organisations and charities, between 1.25 and 3.40 million people in the UK are affected by an eating disorder. The condition can have a tremendous effect not only on a person's mental health and body image but also on his or her oral health, and a patient's mouth can thus serve as a major indicator of disordered eating. In this regard, the Oral Health Foundation has recently highlighted the role of dentists, dental hygienists, dental therapists and dental nurses in recognising the warning signs and symptoms of eating disorders.

According to various studies, eating disorders are among the deadliest mental illnesses. In the UK, around 275,000 people have binge eating disorder, 235,000 bulimia and 100,000 anorexia. Purging through vomiting often leads to enamel erosion, whereas binge eating may increase the risk of dental caries and tooth loss caused by excessive sugar consumption. Additionally, people with eating disorders often absorb insufficient vitamins and other nutrients, and malnourishment too can be reflected in the mouth. Other dental problems related to eating disorders include sensitive teeth, dry mouth, halitosis and enlarged salivary glands.

Dentists have an important role

In light of the National Smile Month campaign aimed at championing the importance of having good oral health, the Oral Health Foundation has recently stated that dental professionals are well placed to identify eating disorders early during regular dental check-ups.

Brooke Sharp, a clinical advice coordinator at Beat, an eating disorder charity based in the UK, stated that eating disorders often go undetected for a long time, since it is difficult to identify the disorder by just looking at a person. However, since there are physical indicators of an eating disorder that are visible in the mouth, she believes that dentists may be among the first to observe these.

As noted by Chief Executive of the Oral Health Foundation Dr Nigel Carter, OBE, it is already the case that dental professionals examine the hard and soft tissue of the mouth and look for signs of tooth erosion during routine dental examinations. They also have the opportunity to spot possible injuries to the mouth from the insertion of foreign objects to cause the person to vomit.

**“The UK needs a far better strategy for diagnosing eating disorders early”
— Dr Nigel Carter, Oral Health Foundation**

Commenting on the issue, Dr Carter said in a press release: “The UK needs a far better strategy for diagnosing eating disorders early. The sooner an eating disorder is suspected or recognised, then the more effective treatment will be.”

Sharp stated that the charity is already taking action to provide dental professionals with the necessary tools for recognising eating disorders and advising patients on the next steps: “At Beat, we are currently working on a webinar series to educate different health sectors about health, which includes dentists, dieticians, and pharmacists. The aim of this training will be to help professionals identify the signs of an eating disorder and understand how to support their patients.”

However, she noted that, although dental teams may encounter patients with eating disorders, it is not their responsibility to treat the condition. Instead, they should direct patients to professional help: “Whilst it's important to increase awareness of eating disorders, it is never the role of dentists to diagnose or treat the eating disorder, and patients must always be signposted to specialist support,” she concluded.

Study indicates denture wearers may be more at risk of nutritional deficiencies

By Anisha Hall Hoppe, Dental Tribune International



A retrospective longitudinal study conducted by the Regenstrief Institute and the Indiana University School of Dentistry indicates that full or even partial denture wearers might be at risk of nutritional deficiencies. (Image: Shutterstock/Krakenimages.com)

Although prosthodontic treatment is the primary means of helping either completely or partially edentulous patients, nutritional counselling too might be warranted for patients who have received dentures, according to a recent retrospective longitudinal study by Regenstrief Institute in Indianapolis and the Indiana University School of Dentistry.

The study linked laboratory values of nutritional biomarkers with dental records. The laboratory tests included complete blood count, basic metabolic profile, and lipid and thyroid panel tests. The research team compared the nutritional state of 3,519 study subjects who had received removable partial or complete conventional or implant- or abutment-retained dentures with the nutritional state of 6,962 members of a control group who had not received or required prosthodontic intervention. In addition to age and sex, the authors considered other factors available in the electronic records, including race, treatment history, and insurance status. The study used records spanning from 1 January 2010 to 31 December 2018 of patients who had been treated at Indiana University School of Dentistry clinics, and nutritional markers over the course of two years were compared.

Noticeable findings included significant decreases in serum albumin, creatinine and blood urea nitrogen (BUN) in completely edentulous patients after they had received their dentures and an increased estimated glomerular filtration rate (eGFR) in the same group. Those who had dentures also had decreased serum albumin, protein and serum calcium compared with controls. For partially edentulous patients, there was a decrease in thyroid-stimulating hormone, BUN and total

cholesterol. Records for patients who were edentulous in either the maxillary or mandibular arch indicated decreased BUN and eGFR as well.

The authors suggested that lowered levels of serum albumin in denture wearers could have a connection to older patients having problems with chewing and swallowing and in patients who already struggle with chronic inflammation and malnutrition. Generally, serum albumin and prealbumin decrease only in patients who experience severe malnutrition, and the study authors recommended further studies to explore the connection between albumin levels and prosthodontic intervention.

Senior author Dr Thankam Thyvalikakath, a research scientist at Regenstrief Institute and director of the dental informatics core at the Indiana University School of Dentistry, shared her perspective in a press release. She said: “Dentures are a significant change for a person. They do not provide the same chewing efficiency, which may alter eating habits. Dentists need to be aware of this and provide advice or a referral for nutrition counselling. These patients need support during the transition and possible continued monitoring.”

The study authors concluded that serological testing could help with monitoring the health and nutritional state of denture wearers and that patients should be offered nutritional counselling when receiving their prosthodontic treatment.

Although serum albumin, calcium, protein and creatinine decreased in denture wearers, the mean values of haematological markers remained within normal ranges, excluding triglyceride, low-density lipoprotein and total cholesterol. Weight and body mass index were above the normal range. The study authors noted that the latter values indicated overweight patients in both the control and study groups and were consistent with the local population in Indiana.

The risk of undernutrition is not a new consideration, but the use of electronic records enabled a much more feasible approach to evaluating the health and progress of a larger selection of patients.

The study, titled “Nutritional assessment of denture wearers using matched electronic dental-health record data”, was published online in the Journal of Prosthodontics on 23 March 2022, ahead of inclusion in an issue.



A new US study has found that the overall median survival time for teeth after undergoing root canal therapy is 11.1 years.
(Image: VanoVasaio/Shutterstock)

Study analyses tooth survival after root canal therapy in US population

By Brendan Day, Dental Tribune International

Though there have been many studies on the survival rate of endodontically treated teeth, only a relatively small proportion of these have analysed these outcomes outside of an academic clinical setting. Using data from a large group of general dental practices across the US, researchers have been able to glean a deeper insight into just how long teeth tend to survive after undergoing root canal therapy and predictors of that survival time.

The study was conducted by a team led by Dr Thankam Thyvalikakath, a research scientist at Regenstrief Institute and director of the dental informatics core at the Indiana University School of Dentistry, both in Indianapolis. The team used de-identified data of 46,702 patients who had received root canal therapy from members of the National Dental Practice-Based Research Network, a consortium of dental practices and organisations throughout the US. According to the researchers, this study is the first of its kind in the US to analyse such records from patients at community dental practices.

Analysis of the data found that the overall median survival time for teeth after undergoing root canal therapy was 11.1 years and that 26% survived for more than 20 years. The researchers also demonstrated that these survival times differed depending on

follow-up treatments and other factors. Teeth that received root canal therapy along with a subsequent filling and crown had a median survival time of 20.1 years, whereas those that received no additional restorative work after the therapy survived for an average of just 6.5 years.

Another significant predictor of tooth survival time after root canal therapy was the presence of dental insurance at any point during dental care. In a press release, Dr Thyvalikakath noted that these findings could “inform dental insurance coverage by demonstrating the value of crowns and permanent restoration options”.

“Oral health is a public health issue that significantly affects people’s overall health. Leveraging dental records can help us better understand ways to improve treatment, identify causal relationships and maintain the health of teeth and gums,” she added.

The study, titled “Root canal treatment survival analysis in National Dental PBRN practices”, was published online in the Journal of Dental Research on 12 May 2022, ahead of inclusion in an issue.



Superior Dental Communication

The most versatile solution for non-interruptive dental communication, documentation and education.

futudent»
SHARE your CARE



microCam XS
When weight matters



proCam XS
When size matters



smartCam
When convenience matters

Save time and effort

Futudent non-interruptive cameras are designed for everyday imaging without interfering with treatment flow.



Futudent makes it easy to take before and after images, case documentation images, and patient education videos in your daily workflow with every patient. This offers tremendous savings in time and excellent tools for growing your business from increased efficiency and treatment acceptance rates to happier patients.

Benefits



- > Easy mounting on chair light or loupes
- > Instant hands-free operation by foot pedal → no treatment interruptions
- > Quick documentation and sharing
- > Simple USB installation and integration
- > Modern case visualization and patient treatment plans
- > Certified and secured products, EU MDR 2017/745 Class I
- > Fast ROI by improvements in treatment acceptance rates
- > Powerful tool for safe and repeatable education
- > Replaces your current interruptive intra-oral and digital cameras

futudent»

For more information about the Futudent non-interruptive camera systems and local distributors, please visit www.futudent.com