implants

international magazine of Oral implantology




case report

The Stable Tissue Concept

events

50 plus one years into the future







Z1[®]Implant

Giving you confidence in implantology.



Proven clinical outcomes for patient safety

- ✓ Anti-bacterial shield
 ✓ Ideal in f Ideal in fresh extraction sockets
 - ✓ Immediate aesthetic result



Proven economics for a profitable implant practice

- ✓ Reduced chairtime
- Practice development
- ✓ Patient satisfaction



Proven surgical protocols for a simplifed workflow

- ✓ Only 1 surgery
 ✓ No healing abutment
- ✓ Visibility of the connection



Proven Technology for a better integration

- ✓ 98.6%* success rate
- ✓ Pure Titanium and Y-TZP Zirconia
- ✓ Suitable for all prosthetic solutions







Dr Rolf Vollmer

First Vice President and Treasurer of DGZI

50 years of DGZI—a strong indicator for European dental implantology

In celebration of the 50-year anniversary of the German Association of Dental Implantology (DGZI), its third Future Congress for Dental Implantology was held in Cologne in Germany in early October. Owing to coronavirus-related travel and other restrictions imposed by governments worldwide, we had to celebrate our anniversary with a delay of one year, but this did not hinder its success in any way. We can proudly look back on a beautiful and worthy celebration of half a century of European implantology and a multifaceted, exciting congress. All in all, this one-of-a-kind event did full justice to the unique occasion.

As the oldest implantological expert society in Europe, celebrating 50 years of our existence was truly a milestone for us. We used the special event in Cologne as an occasion to reflect on the incredible developments that dental implantology has undergone in the past 50 years since our foundation by the visionary group led by Prof. Hans Grafelmann. DGZI has accompanied and helped to shape these developments from their very beginnings until today. Moreover, through our anniversary congress, we have succeeded in providing a visionary outlook of what future implantology might offer in five to ten years from now in terms of new clinical techniques and new approaches to implantology in general.

We can proudly say that our valued guest speakers were indeed the who's who of European implantology and

contributed greatly to the scientific programme. Against this background, I would like to express my most sincere gratitude to the presidents and board members of the German Society for Implantology (DGI), the German Society of Oral Implantology (DGOI) and the other specialist societies who accepted our invitation to address our congress participants and delivered truly insightful lectures. Despite the undoubtedly competitive situation between the various professional associations, the 2021 event in Cologne revealed something quite important: on certain topics and in certain situations, those at the forefront of German—and on a broader scale of course European—dental implantology are united in their vision of the future orientation of implantology. This is a strong indicator for us!

In this spirit, I would like to extend warm and friendly greetings to you and wish you enjoyable reading of the last issue of this anniversary year of implants-implantology, as well as a reflective time ahead of the Christmas season!

Sincerely yours,

a loss an

Dr Rolf Vollmer







Cover image courtesy of Argon Dental / www.argon-dental.de



editorial

50 years of DGZI—a strong indicator for European dental implantology O3 Dr Rolf Vollmer

research

The Implant Protection Plan (I.P.P.) 06 Drs Tiziano Testori, Giordano Bordini & Matteo Basso The role of metallic nano- and microparticles in peri-implantitis 14 Dr Ioannis Papadimitriou

case report

The Stable Tissue Concept 20 Dr Kai Zwanzig Sinus augmentation and simultaneous implant placement 24 Drs Fernando Duarte, Carina Ramos, Paulo Veiga & Marco Infante da Câmara Alveolar deficiency management in maxillary lateral incisor agenesis Dr Federico Berton Immediate functional implants in the aesthetic zone of a heavy smoker 32 Dr Dr Branislav Fatori & Dr Inge Schmitz

interview

Fine dentistry and creative engineering go hand in hand 34 An interview with Dr Kai Zwanzig and Ric Donaca

news

manufacturer news 38 news 48

events

50 plus one years into the future 42 Dr Georg Bach

about the publisher

50 imprint



NovoMatrix™ Reconstructive Tissue Matrix the next generation material

NovoMatrix™ Reconstructive Tissue Matrix is an acellular dermal matrix derived from porcine tissue intended for soft tissue applications. The proprietary LifeCell™ tissue processing is designed to maintain the biomechanical integrity of the tissue, which is critical to support tissue regeneration.

Indications

- Localized gingival augmentation to increase keratinized tissue (KT) around teeth and implants
- Alveolar ridge reconstruction for prosthetic treatment
- Guided tissue regeneration procedures in recession defects for root coverage

MinerOss™ A is a trademark of BioHorizons®. BioHorizons® is a registered trademark of BioHorizons.

©BioHorizons. All rights reserved. Not all products are available in all countries

www.biohorizonscamlog.com

Product features

- Consistent thickness (1 mm)
- Pre-hydrated
- Controlled source

Before use, physicians should review all risk information, which can be found in the Instructions for Use attached to the packaging of each NovoMatix™ Reconstructive Tissue Matrix graft. NovoMatrix™ is a trademark of LifeCell™ Corporation, an Allergan affiliate. ©BioHorizons. All rights reserved Not all products are available in all countries.





The Implant Protection Plan (I.P.P.)

Innovative implant and periodontal maintenance protocol

Drs Tiziano Testori, Giordano Bordini & Matteo Basso, Italy

Introduction

The last 30 years of world dentistry have undoubtedly been characterised by the exponential growth of implantology, which has gone from being a discipline in the hands of a few experts "to being a field of treatment at many dental practices. There are several reasons for this increase: firstly, surgical and prosthetic techniques have been simplified over the years, repeatable protocols being certified by decades of literature, reducing costs for the patient and limiting invasiveness and postoperative discomfort. In addition, many patients want an aesthetic and functional restoration by means of fixed implant prostheses as their first choice, rather than resorting to solutions such as removable prostheses or fixed prostheses on natural teeth involving the prosthetic

"... modern implantology is trying to address the possibility of guaranteeing a clinical result that endures over time ..."

preparation of healthy teeth. Ultimately, it should not be under-estimated that implantology has also increased because it represents a source of income for the economic balance sheet of many healthcare facilities that have decided to specialise in this field. However, the implementation of an oral implant rehabilitation, be it a single tooth or a complex solution, cannot and must not today represent the end point either for the patient or for the dentist and his or her team. Nowadays, thanks to our knowledge, we have no difficulty in achieving implant-based rehabilitation even in cases of severe bone atrophy using regenerative techniques.

The critical point that modern implantology is trying to address, not always successfully, is the possibility of guaranteeing a clinical result that endures over time.

To achieve this ambitious goal, it is crucial to design an effective and feasible implant and periodontal maintenance protocol. We know that home maintenance around implants can be more difficult than around natural teeth because the techniques and instruments to be used, in many clinical cases, are inevitably different from those used for natural teeth. In addition, we may be confronted with the typical pathologies of implants, represented by mucositis and peri-implantitis, subtle pathologies that are difficult to control and whose differences from gingivitis and periodontitis we have learnt about.1 According to studies on the prevalence of peri-implant disease,² 45% of patients show, after an average of nine years, signs of mild periimplantitis and 14.5% medium to severe. In recent times, we have gained knowledge about oral biofilm, discovering that the biofilm changes in its characteristics when a pathology is established and that some pathologies, such as mucositis and peri-implantitis, are characterised by a repetitiveness in the type of pathogenic microorganisms present.3 Implant maintenance protocols, however, have not evolved alongside knowledge, sometimes only introducing new instruments or technologies, such as laser therapy or phototherapy, and some new antiseptic principles.

The concepts of periodontal and peri-implant eubiosis and dysbiosis

One of the key points for the long-term success of a patient rehabilitated with implants, which is no different from that of a patient treated for periodontal disease, is to establish a correct programme of supportive therapy and periodic follow-up that includes differentiated recalls based on an analysis of risk factors and consequent classification into risk categories. The literature and our decades of clinical experience have shown that patients with treated periodontal disease are at risk of having setbacks and developing a new disease process.4 Thus, the implant patient or the periodontal therapy patient should not and must not be considered a patient who after treatment, however successful, can return to being normal and be low risk. Based on this scientific and clinical evidence, we can begin to plan the future of our therapies, starting with the biological basis of the problem and the new assump-



AN EFFECTIVE PREVENTION against periodontal risk factors





Dental implants



Natural teeth treated for previous parodontosis





Smokers



Reduced immune defences

OZONATED OLIVE OIL, PVP/VA, COLOSTRUM AND STEVIA

B. LACTIS HN019 - MARXIANUS FRAGILIS - COLOSTRUM - BIOTIN



Curasept Prevent is an innovative line designed, for daily use, to mantain a balanced oral microbiota. It is ideal in the management of prevention protocols of mucositis and peri-implantitis when there are dental implants. Moreover it can be useful in the prevention of gingivitis in periodontal patients, with natural teeth, and in presence of particular risk factors that can facilitate the onset of oral diseases.






Fig. 1: Operational checklist for maintenance sessions.

tions linked to a more accurate knowledge of the oral microbiota.

The oral microbiota is the set of microorganisms that live and coexist in the oral cavity. It should be distinguished from the concept of oral microbiome, which is the collective genomes of the microorganisms present. The microbiota is made up of more than 700 different bacterial species, as well as numerous other microorganisms, and in a healthy state it is in perfect balance with the host, causing no harm and providing numerous benefits through the predigestion of food, antibacterial action, and the secretion of enzymes. This state of equilibrium is called "eubiosis". It is important to emphasise that a eubiotic microbiota may also contain species that are considered pathogenic but which as part of a balanced biofilm are not capable of inducing pathology. Thus, the mere presence of periodontopathogenic species or implant pathogens is no longer considered a sign of pathology, as it was in the past; they can at most be considered risk factors to which more attention should be paid.

However, when a pathological process of either periodontitis or peri-implantitis occurs, the oral biofilm changes and a picture of dysbiosis⁵ of the oral microbiota emerges, there being a change in the relationships between the present species and that can trigger an immune and inflammatory response. It is precisely the inflammation created by the bacterial trigger that feeds and often maintains the dysbiosis itself, leading to chronicity of the condition. This alteration in the balance leading to the onset of disease is affected by many variables or risk factors which can affect the patient's clinical situation at several levels. 6 There is a solid body of literature on the most important risk factors for periodontal disease, drawn from many clinical trials and a smaller number of longitudinal studies. 7,8 This has made it possible to identify some of these factors as being strongly correlated with periodontal disease and, at least regarding the current state of research, to suggest for others a correlation whose nature has yet to be validated in detail.9-14

Risk factors include some that are modifiable and others that are not. Among the most important modifiable factors are smoking, stress and diabetes, which we know cannot yet be eliminated but is treatable and therefore modifiable. Among the non-modifiable factors is genetic predisposition, a generic and imprecise term that refers to a host's ability to modulate the quality of the immune and inflammatory response differently and thus favour the onset of disease. Other risk indicators whose correlation with implant and periodontal disease has yet to be fully clarified include obesity, metabolic syndrome, hypertension, cardiovascular disease, and osteopenia/ osteoporosis.

A cooperative patient-professional plan

Once the biological bases currently considered valid for a more scientific and modern understanding of periimplant pathologies have been clarified, other much more practical and organisational aspects of maintenance need to be investigated. First of all, not all patients are the same: the selection of a good candidate for periodontal treatment or implant therapy should always be made a priori, excluding those patients in whom the risk factors described, or even who display an unsuitable propensity and attitude regarding adhering to the practitioner's requests and prescriptions (patients defined as having a low degree of cooperation or compliance), are not considered satisfactory. Sometimes it is not possible to select only ideal candidates, and even these, in the course of their lives, may suffer a disease setback if not properly motivated and followed up. Although a number of periodontal and peri-implant risk assessment tools have existed for years to assist clinicians in setting up the most suitable maintenance programme possible, 15,16 there is no uniformity even among the most experienced professionals.¹⁷ Furthermore, the patient undergoing implant and periodontal rehabilitation often requires a guarantee of duration of treatment, a need which goes far beyond what medicine can offer but to which, nowadays, we cannot fail to provide a satisfactory answer.

The aim of this scientific contribution is to suggest a protocol, the implant protection plan (IPP), that establishes a therapeutic alliance between the treating dentist, the dental hygienist and the patient at the end of the active phase of periodontal or implant therapy. The IPP protocol provides for a shared maintenance pathway which starts with the initial assessment of the patient and periodic re-evaluations, which are not an end in themselves or a mere collection of clinical data, but determine actions and changes in the dental professional's attitude or that of the patient in order to optimise the periodontal and implant prognosis. The patient himself or herself should feel involved in the IPP, share its purpose and not play a passive role (Figs. 1 & 2).

The protocol also includes an operational checklist with all the factors that the hygienist has to check during the session. This tool is designed to monitor the clinical situation and alert the dental practitioner to any worsening of the patient's clinical condition compared with the baseline and to make consequent adjustments to the current supportive therapy (e.g. shortening recall times; Fig. 3).

The true innovation is to ensure that the patient does not have to give up compliance because, in return for a personalised maintenance programme set out in a contract signed by both parties, he or she will be guaranteed specific treatments or interventions, such as prosthetic replacement treatments, without any financial cost should any biological problems occur, but only if he or she has complied with the maintenance sessions agreed with him or her beforehand.

The first step in implementing the protocol is to assign a periodontal or peri-implant risk profile. The dentist determines a specific risk class on the basis of systemic and local risk factors, the presence or absence of implants (patient with only natural teeth, patient with natural teeth and implants, or patient with only implants) in order to plan the frequency and manner of individualised maintenance therapy. The assessment of the risk profile is therefore divided into a periodontal profile, if the patient still has natural teeth, and a periodontal framework in order to combine the two classifications into a single patient risk class assignment. Assigning a prognostic risk for a patient who has a partial natural dentition is a process that requires cross-referencing anamnestic information and elements from the objective examination and interpreting this data through prognostic assessment. It is necessary to include in the

ame a	nd Surname		PPORTIV	ETHERAP	Y PLANNING
INITIAL ASSESSMENT RISK CATEGORY			1-YEAR ASSESSMENT RISK CATEGORY		
□ s	TAGE II B/C	TYPE 1		STAGE II B/C	TYPE 1
□ s	III A	TYPE 2		STAGE IV A	TYPE 2
□ s	III D/C	TYPE 3		STAGE III B/C	TYPE 3
	FOLLOW-UP	PATIENT COMPLIANCE		FOLLOW-UP	PATIENT COMPLIANCE
YPE 0	6 months 4 months	☐ Medium/High	TYPE 0	6 months 4 months	☐ Medium/High ☐ Low
YPE 1	4 months 6 months	Low Medium High	TYPE 1	4 months 6 months	☐ Low ☐ Medium ☐ High
YPE 2	3 months 4 months	Low Medium High	TYPE 2	3 months 4 months	☐ Low ☐ Medium ☐ High
YPE 3	2 months 3 months	Low Medium High	TYPE 3	2 months 3 months	☐ Low ☐ Medium ☐ High

Fig. 2: Programming of periodontal and implant maintenance sessions.

analysis the patient's medical and dental history, oral and extra-oral radiographs and the main periodontal variables (plaque index, bleeding on probing, probing depth, recessions, furcation involvement, pathological tooth mobility, bone profile) and to give the patient the correct periodontal disease diagnosis (in terms of stage and grade).¹⁸

Today, there are several tools and algorithms that help us to plan the correct timing of maintenance sessions and, indirectly, to predict the patient's prognosis. It should also be pointed out that some of these collected variables, besides having a greater relative weight (odds ratio) than others in influencing prognosis, offer more information as indicators of disease progression. The most important of these are smoking, diabetes and a history of periodontitis.

In a recent review of the scientific evidence supporting periodontal maintenance planning, the following was emphasised:¹⁹