ortho

international magazine of Orthodontics

Are you ready for **E-xcellere** DIGITAL | 2020

3M ORTHO CONFERENCE

please visit

www.excellere3m.com

research

Auto-administered photobiomodulation: A new way to make orthodontic treatment faster and more comfortable?

opinion

Accelerated aligners with photobiomodulation

feature


How COVID-19 has forced us all to relook our practices, lives and goals



DIGITAL INNOVATIONS FORUM FUTURE PROOFING YOUR PRACTICE BY EMBRACING TECHNOLOGY

For dental professionals seeking a competitive edge through digital innovation.

Align Technology will soon host its first ever online Digital Innovations Forum. This 2-day event will showcase the latest digital innovations and demonstrate how they can help provide improved and more efficient treatment experiences for your practice, and your patients.







Dr Sean K. Carlson

CEO of OrthoScience

Why caring matters more than ever

The world of technology is advancing quickly. Before mobile devices and smartphones, communication was not instantaneous. Whether you had news or just a simple greeting to convey, in the old days, you could not reach people quickly, nor could they reach you. This change is important. It affects healthcare in a big way. It affects patient expectations and it affects doctor efficiency. Never has this been clearer than with the current climate brought about by COVID-19. We have all become instantly proficient at digital and remote communication—instant communication. It is inevitable that orthodontists will be expected to increase instant accessibility and remote solutions more than ever before. However, despite what may sound like added convenience, the new world of remote communication and instant access is filled with pitfalls that will cause many doctors to stumble in this new world.

Pitfall 1: Easy access does not mean automation

You cannot phone in caring. Computer bots that provide automated responses will become frustrating for patients. Healthcare is about caring. Patients go to doctors for care, not products. When we forget this basic principle, we are no longer doing our job. An automated system may work for your cellular service provider, but it will not work for your doctor. Orthodontists should find digital communication solutions that are personal. Patients demand and deserve a personal touch. They deserve caring. Keep this in mind when seeking out your future digital solutions.

Pitfall 2: Patients are not customers

Patients are people with a disease. They are not shopping for a new hat. Although orthodontic disease is rarely, if ever, life-threatening, malocclusions still demand the attention that only a trained doctor can provide. And just like the first pitfall, you cannot phone in or automate treatment. This has never been clearer than with the experiment of automated orthodontics that is being attempted by so many direct-to-consumer tooth alignment products. These are emerging with increasing speed in the US. Without doctor supervision, the failures and shortcomings of these approaches simply cannot be appropriately managed. Patients are the ones who end up

suffering. Unfortunately, since the shortcomings of these do-it-yourself orthodontic solutions are rarely life-threatening, the consequences of negative outcomes will be overlooked for a long time to come. But practising orthodontists understand that finishing orthodontic cases requires careful attention. It is not easy. When patients are treated like customers, you only need to sell the start. When patients are treated like patients, you are fully aware that you need to provide the finish—the cure to the malocclusion. This requires a much greater commitment on the doctor's part.

Pitfall 3: Thinking that technology will make caring easier

Unfortunately, the job of a doctor is difficult. Caring about your patients is difficult. Technology will not eliminate this. Making patients well—curing their disease—is burdensome. Just because some things become automated does not mean that you can offload this burden. However, technology will allow you to care better, that is, your connection to your patients will be stronger than ever. You will have the ability to control the direction of your treatment on a micro-scale. This is where technology will shine. Traditionally, doctors would connect with patients only at physical appointments. Now they can connect with them remotely far more often. This increased connection should not be viewed as added work: it should be viewed as added control. For the doctor, this means fewer physical visits, more care opportunities, better appliance monitoring, fewer phone calls, more patient touch points. All of these things, when viewed through a caring lens, will improve our practices tremendously. So how will you embrace new technology? When making your own choices, I encourage you to view your smartphone through the eyes of a patient. Imagine you have a disease and you have just employed a team of doctors to cure you. How could your smartphone help you? How would your doctor team show you that they care about you through your smartphone? How would they let you know they care about your healing and your cure just as much as you do? Once you have envisioned that scenario, you will understand clearly what your technology solutions need to encapsulate. I trust that the thing you will want to feel the most is that the doctors care.







Cover image: **3M** www.3M.com



)4| ortho


international imprint

editorial Dr Sean K. Carlson	03
research	
Auto-administered photobiomodulation: A new way to make orthodontic treatment faster and more comfortable? Drs Carlo Fornaini & Elisabetta Merigo	06
technique New Age orthodontics and orthopaedics with temporary anchorage devices: Unlocking our clinical potential Drs Trevor S. Nichols & Stuart L. Frost	10
case report Is orthodontic treatment compatible with a healthy periodontium? A narrative review Drs Francisco Molina & Carlos A. Molina	22
clinical case Successful treatment outcomes using 3M Clarity Ultra Self-Ligating Brackets bonded using a digital flash-free bonding technique Dr Riccardo Riatti	26
opinion	
Accelerated aligners with photobiomodulation Dr Miguel Stanley	34
Artificial intelligence shifts consumer behaviour and impacts dentistry Dr Derek Mahony	40
feature How COVID-19 has forced us all to relook our practices, lives and goals Jerko Bozikovic	48
Will COVID-19 push orthodontics further into the digital space? Jeremy Booth	52
interview In-office aligners: Taking control of orthodontic laboratory expenses Dr Melissa Shotell	54
Orthodontists should see their patients through the eyes of a periodontist Dr Jean-Marc Dersot	58
business news Clear aligner treatment models jostle for acceptance in the pandemic Jeremy Booth	60
Clinical director of CA DIGITAL joins inviSolution InviSolution	62
events First digital Invisalign Scientific Symposium: Scientific data, shared experiences, peer recognition, social responsibility	64
about the publisher submission guidelines	68

70

MRESTRO 3D

www.maestro3d.com



2 x 5.0 MEGAPIXEL


Plug & Play SS

Smart Impression Scanning

Accuracy less than 8 micron

Texture Superimposition


Study Models



Clear Aligner



Virtual Setup



Indirect Bonding



Auto-administered photobiomodulation: A new way to make orthodontic treatment faster and more comfortable?

Drs Carlo Fornaini & Elisabetta Merigo, Italy

"Photo-biomodulation (PBM) therapy" is the universally recognised term for the wide range of laser applications with low parameters, replacing the terms of "low-level laser therapy", "biostimulation", "soft laser" or "cold laser", mostly used in the 1970s and 1980s.¹-³ This term was added in 2015 to the National Library of Medicine's controlled vocabulary thesaurus, Medical Subject Headings.⁴

The father of PBM is the Hungarian physician Endre Mester, who first observed in the 1960s the effects of a ruby laser (wavelength of 694.3 nm) he used on animal models in order to seek to understand any carcinogenic effect of laser used at low energy densities (1 J/cm²).⁵ Mester performed studies on the effect of laser phototherapy on healing processes and tissue repair in animals, on phagocytosis of bacteria by leucocytes, on synthesis of haemoglobin and on healing of ulcerative lesions non-responsive to conventional therapies.⁶

The scientific interest around PBM is due to its various properties in terms of stimulation of both wound healing (of mucosa, skin, tendon, etc.) and repair (bone, cartilage, dentine), as well as pain and inflammation.

On the basis of its effects, PBM is defined by Anders et al. as "a form of light therapy that utilises non-ionizing forms of light": it is a photothermal reaction which involves endogenous chromophores resulting in beneficial therapeutic phenomena, the most important being the alleviation of pain or inflammation, immunomodulation, and promotion of wound healing and tissue regeneration".

The photochemical reactions at the base of PBM can produce three different clinical results: stimulation of healing, an anti-inflammatory effects and analgesia.^{5, 6} *In vitro* and *in vivo* studies on animals and humans have demonstrated the efficacy of PBM in promoting



Fig. 1: The laser device for auto-administered photobiomodulation.

physiological effects such as DNA synthesis, neoangiogenesis, keratinocyte proliferation, fibroblasts and endothelial cells proliferation, collagen synthesis and deposition, activation of macrophages, revascularisation and contraction of the wound.^{7,8}

PBM may increase the activity of macrophages and neutrophils and has a specific and preferential modality for certain mediators of inflammation. PBM inhibits the catabolic mediators of inflammation, such as inhibitors of collagen synthesis and cell proliferation; reduces the influx

of neutrophils to the level of inflamed tissue and stimulates the production of anti-inflammatory metabolites such as cyclo-oxygenase 1 (COX-1) and 2 (COX-2). PBM also seems to contribute to the reduction of oedema. The mechanism underlying the analgesic effect of PBM has not yet been fully clarified and is probably quite complex: one of the most accredited hypotheses is related to the increase of the nociception threshold with a neural block, specifically with inhibition of A delta and C nerve fibres mediated by alteration of axonal flow or by inhibition of neural enzymes. Furthermore, there appears to be an increase in the production of endorphins due to changes in opioid receptors. PBM can also mimic the effects of anti-inflammatory drugs by attenuating the level of prostaglandin E₂ and by inhibiting COX-2.10

PBM results in the reduction of acute and chronic pain through a conduction block and alteration of A delta and C nociceptive nerve fibres, action occurring at the level of the central nervous system through ascending and descending transmission. Primary reactions to PBM occur in the irradiation zone, but a secondary systemic response related to the transport of photoproducts such as prostaglandins, enkephalins and endorphins mediated by the lymphatic system and with a persistent effect for several hours or weeks is thought possible. This is the main reason why it is important in the design of experimental studies and in the interpretation of the results to consider the use of protocols in which there is an internal control. Proceedings of the state of the second of the results to consider the use of protocols in which there is an internal control.

Thanks to the use of parameters characterised by low energy density, PBM can be considered free of possible side effects compared with higher laser powers correlated to damage and tissue destruction.^{12, 13}

The application of PBM in the orthodontic field has seen in recent years increased scientific production, initially in in vivo studies in animals and subsequently in clinical trials aimed at investigating the effect of the application of laser protocols at low energy on the acceleration of tooth movement with a simultaneous analgesic potential.14 The stimulatory effect of the proliferation and differentiation of the cells involved in bone metabolism, in particular osteoblasts and osteoclasts, is the basis of the results obtained. At the maxillary and mandibular level, this results in the reduction of the time needed to finalise orthodontic therapy. This type of application, free from side effects, especially regarding the periodontium, defines a beneficial effect in the prevention of carious disease and a simultaneous improvement of the pain symptoms often related to the active phases of orthodontic movement.15

Systematic reviews and meta-analyses of existing literature on PBM and the acceleration of tooth movement in the Cochrane Library, PubMed Central, Embase, Scopus



Fig. 2: Auto-administered photobiomodulation for the treatment of temporomandibular disorder.




Fig. 3: Auto-administered photobiomodulation just after orthodontic bonding.

and Web of Science showed a statistically significant difference in the acceleration of tooth movement between treatment with PBM and non-laser treatment.^{16, 17} Several *in vivo* randomised controlled clinical trials evidenced that

photobiomodulation is effective in alleviating the intensity and duration of pain experienced by patients after the engagement of alignment archwire.^{18, 19}

Studies in the literature, in particular, on protocols at different times of administration of PBM (weekly, every other day, with monthly precise doses) report more striking effects for low fluence values (between 5 J/cm² and 8 J/cm²) compared with values above 20 J/cm².

Unfortunately, a limiting factor of this kind of treatment is that the patient needs to attend therapy at least two or three times a week for treatment, and this may negatively influence patient compliance. To allow the patient to receive treatment without visiting the therapist, there recently appeared on the market a new family of devices that, owing to their Class II classification by the American National Standards Institute, may be used directly by the patient without the need for protective goggles in a very simple way, thanks to the preset parameters. Device size and cost are reduced and contra-indications and the risk of side effects are absent; nevertheless, evaluation of the patient by a specialist is mandatory (Figs. 1–3).

At-home laser use has been described for temporomandibular disorder, for example, using preset devices and therapies determined by the therapist, limiting the discomfort of repeated appointments and obtaining good results in terms of pain.²³ Some case reports have also described laser use for the treatment of neurological face diseases, also provoked by a wrong insertion of intraosseous implant for prosthetic rehabilitation,²⁴ as well as diabetic leg vascular lesions treatment.^{24, 25}

We think that orthodontists might today consider these new opportunities as an easy way to minimise the treatment time and to reduce or eliminate pain during the first step of the therapy, thereby increasing patient compliance.

about



Dr Carlo Fornaini is a researcher at the MICORALIS laboratory of the dental faculty of the Université Côte d'Azur in Nice in France. He has a private practice in Fiorenzuola d'Arda in Italy.



Dr Elisabetta Merigo is also a researcher at the MICORALIS laboratory.

References:

- 1 Enwemeka CS. Low level laser therapy is not low. Photomed Laser Surg. 2005 Dec;23(6):529–30. doi: 10.1089/pho.2005.23.529.
- 2 Karu T. Photobiology of low-power laser effects. Health Phys. 1989 May;56(5):691–704. doi: 10.1097/00004032-198905000-00015.
- 3 Reddy GK. Photobiological basis and clinical role of low-intensity lasers in biology and medicine. J Clin Laser Med Surg. 2004 Apr;22(2):141–50. doi: 10.1089/104454704774076208.
- 4 Anders JJ, Lanzafame RJ, Arany PR. Low-level light/laser therapy versus photobiomodulation therapy. Photomed Laser Surg. 2015 Apr;33(4):183–4. doi: 10.1089/pho.2015.9848.
- Mester E, Spiry T, Szende B, Tota JG. Effect of laser rays on wound healing. Am J Surg. 1971 Oct; 122(4):532–5. doi: 10.1016/0002-9610(71)90482-x.
- Tunér J, Hode L. Laser therapy in dentistry and medicine: a guide for research scientists, dentists, nurses, dental hygienists, and other interested parties within the medical field. Grängesberg: Prima Books; 1996. 233 p.
- 7 Mester E, Mester AF, Mester A. The biomedical effects of laser application. Lasers Surg Med. 1985;5(1):31–9. doi: 10.1002/lsm.1900050105.
- 8 AlGhamdi KM, Kumar A, Moussa NA. Low-level laser therapy: a useful technique for enhancing the proliferation of various cultured cells. Lasers Med Sci. 2012 Jan;27(1):237–49. doi: 10.1007/s10103-011-0885-2.
- 9 Kingsley JD, Demchak T, Mathis R. Low-level laser therapy as a treatment for chronic pain. Front Physiol. 2014 Aug 19;5:306. doi: 10.3389/fphys.2014.00306.
- 10 Chagas LR, Silva JA Jr, de Almeida Pires J, Costa MS. Expression of mPGES-1 and IP mRNA is reduced by LLLT in both subplantar and brain tissues in the model of peripheral inflammation induced by carrageenan. Lasers Med Sci. 2015 Jan;30(1):83–8. doi: 10.1007/s10103-014-1622-4.
- 11 Kahraman SA. Low-level laser therapy in oral and maxillofacial surgery. Oral Maxillofac Surg Clin North Am. 2004 May;16(2):277–88. doi: 10.1016/j. coms.2004.02.002.
- 12 Sun G, Tunér J. Low-level laser therapy in dentistry. Dent Clin North Am. 2004 Oct;48(4):1061–76, viii. doi: 10.1016/j.cden.2004.05.004.
- 13 Navratil L, Kymplova J. Contraindications in non-invasive laser therapy: truth and fiction. J Clin Laser Med Surg. 2002 Dec;20(6):341–3. doi: 10.1089/104454702320901134.
- 14 Ge MK, He WL, Chen J, Wen C, Yin X, Hu ZA, Liu ZP, Zou SJ. Efficacy of low-level laser therapy for accelerating tooth movement during orthodontic treatment: a systematic review and meta-analysis. Lasers Med Sci. 2015 Jul;30(5):1609–18. doi: 10.1007/s10103-014-1538-z.

- 15 Almallah MM, Almahdi WH, Hajeer MY. Evaluation of low level laser therapy on pain perception following orthodontic elastomeric separation: a randomized controlled trial. J Clin Diagn Res. 2016 Nov;10(11):ZC23–ZC28. doi: 10.7860/JCDR/2016/ 22813.8804.
- 16 AlShahrani I, Togoo RA, Hosmani J, Alhaizaey A. Photobiomodulation in acceleration of orthodontic tooth movement: a systematic review and meta analysis. Complement Ther Med. 2019 Dec;47: 102220. doi: 10.1016/j.ctim.2019.102220.
- 17 Cronshaw M, Parker S, Anagnostaki E, Lynch E. Systematic review of orthodontic treatment management with photobiomodulation therapy. Photobiomodul Photomed Laser Surg. 2019 Dec;37(12):862–8. doi: 10.1089/photob.2019.4702.
- 18 Lo Giudice A, Nucera R, Perillo L, Paiusco A, Caccianiga G. Is low-level laser therapy an effective method to alleviate pain induced by active orthodontic alignment archwire? A randomized clinical trial. J Evid Based Dent Pract. 2019 Mar;19(1):71–8. doi: 10.1016/j.jebdp.2018.11.001.
- 19 Matys J, Jaszczak E, Flieger R, Kostrzewska-Kaminiarz K, Grzech-Leśniak K, Dominiak M. Effect of ozone and diode laser (635 nm) in reducing orthodontic pain in the maxillary arch–a randomized clinical controlled trial. Lasers Med Sci. 2020 Mar;35(2):487–96. doi: 10.1007/s10103-019-02896-0.
- 20 Gavish L, Houreld NN. Therapeutic efficacy of home-use photobiomodulation devices: a systematic literature review. Photobiomodul Photomed Laser Surg. 2019 Jan;37(1):4–16. doi: 10.1089/photob.2018.4512.
- 21 Juhász ML, Levin MK, Marmur ES. A review of available laser and intense light source home devices: a dermatologist's perspective. J Cosmet Dermatol. 2017 Dec;16(4):438-43. doi: 10.1111/jocd.12371.
- 22 Fornaini C, Pelosi A, Queirolo V, Vescovi P, Merigo E. The "at-home LLLT" in temporo-mandibular disorders pain control: a pilot study. Laser Ther. 2015 Mar 31;24(1):47–52. doi: 10.5978/islsm.15-OR-06.
- 23 Merigo E, Rocca JP, Oppici A, Cella L, Fornaini C. At-home laser treatment of oral neuronal disorders: case reports. J Clin Exp Dent. 2017 Apr 1;9(4):e595–e598. doi: 10.4317/jced.53373.
- 24 Raizman R, Gavish L. At-home self-applied photobiomodulation device for the treatment of diabetic foot ulcers in adults with Type 2 diabetes: report of 4 cases. Can J Diabetes. 2020 Jul;44(5): 375–8. doi: 10.1016/j.jcjd.2020.01.010. Epub 2020 Feb 4.
- 25 Merigo E, Tan L, Zhao Z, Rocca JP, Fornaini C. Auto-administered photobiomodulation on diabetic leg ulcers treatment: a new way to manage it? Case Rep Med. 2020 May 20;2020:7428472. doi: 10.1155/2020/7428472.