

laser

international magazine of laser dentistry



I | S | L | D

research

Effects of 10,600 nm carbon dioxide lasers on preventing caries

case report

Complex management of drug-induced gingival hyperplasia

interview

A double-edged sword



SkyPulse®

A new generation of compact
Fotona lasers

A compact, easy
to use Er:YAG Laser
tailored for Your
practice!



Fotona[®]
choose perfection

Dr Stefan Grümer

Treasurer of DGL and ISLD



Software **innovation** at its best

Dear laser friends and colleagues,

We all believe in a bright future for laser dentistry, a specialist field that is guided by experience and driven by innovation. It can be argued that advancements in laser dentistry should be made not only in terms of hardware, but also in terms of software.

Against this background, I would like to cordially invite you to join us for the upcoming annual meeting of the German Association for Laser Dentistry (Deutsche Gesellschaft für Laserzahnheilkunde—DGL), which will be held on 6 and 7 November 2020 in Bremen in Germany. Packed with scientific lectures from highly esteemed colleagues and technological innovations from the industry, the conference is considered a definite highlight by many who regularly seek out renowned dental training events. This 2020 annual meeting will be held as joint event together with the 3rd Future Congress of the German Association of Dental Implantology (Deutsche Gesellschaft für Zahnärztliche Implantologie—DGZI). In this way there will be ample opportunity for extensive interdisciplinary

exchange with colleagues through shared lectures, poster presentations and industry exhibits.

With so-called “table clinics” the programme will feature yet another novelty on the first congress day on Friday: during round table presentations led by experienced trainers, small groups of participants will have the opportunity to discuss various topics revolving around laser dentistry and oral implantology with a view to synergetic effects between these two specialist fields. The participants will also be allowed to switch between tables, allowing them to get a glimpse into many of the topics discussed. The conference will put particular focus on practice-oriented aspects, which laser enthusiasts can implement effortlessly and rapidly into their daily clinical work.

With this in mind, I welcome you to Bremen this autumn and I wish you a great and interesting read with this first 2020 issue of *laser—international magazine of laser dentistry*!

Dr Stefan Grümer



Cover image:
© James Thew – stock.adobe.com



editorial

Software **innovation** at its best
Dr Stefan Grümer

03

research

Effects of 10,600 nm **carbon dioxide** lasers on preventing caries
Kenneth Luk, Ollie Yiru Yu, May Lei Mei, Norbert Gutknecht,
Chun Hung Chu & Irene Shuping Zhao

06

case report

Treatment of midline **diastema**
Dr Nataša Prebil

14

Complex management of **drug-induced** gingival hyperplasia
Dr Mihail Tanev, Dr Georgi T. Tomov & Jyuhn H. Ke

18

Marsupialisation of an oral **mucocoele** with a diode laser
Dr Maziar Mir, Dr Masoud Mojahedi, Dr Mina Mazandrani,
Aysan Shabani & Dr Masoud Shabani

22

industry

Are you a laser dentist?
Dr Dimitris Strakas

26

practice management

Successful **communication** in your **daily practice**
Dr Anna Maria Yiannikos

28

Feedback done **right**
Andrea Stix, MSc, MBA

30

interview

A double-**edged** sword

32

events

Master of Science course "**Lasers in Dentistry**" offered again
2020 **annual congress** of DGL to take place in Bremen
Prof. Andreas Braun

34

35

news

manufacturer news

36

news

37

about the publisher

imprint

42



M.Sc. Lasers in Dentistry

Specialist in dental laser therapies

Next Start: **17th September 2020** | Aachen, Germany | 4 semesters

Aachen Dental Laser Center - AALZ
Pauwelsstraße 17 | 52074 Aachen | Germany
phone +49 241 47 57 13 10 | fax +49 241 47 57 13 29
info@aalz.de
www.aalz.de

RWTH International Academy
Kackertstraße 10 | 52072 Aachen | Germany
phone +49 241 80 23543 | fax +49 241 80 92525
info@academy.rwth-aachen.de
www.academy.rwth-aachen.de



Aachen Dental Laser Center

**INTERNATIONAL
ACADEMY**

**RWTH AACHEN
UNIVERSITY**

PROFESSIONAL EDUCATION PROGRAMMES

Your contact for more information: Leon Vanweersch • vanweersch@aalz.de - www.aalz.de

Effects of 10,600 nm carbon dioxide lasers on preventing caries

A literature review

Kenneth Luk^{1,2}, Ollie Yiru Yu², May Lei Mei², Norbert Gutknecht³, Chun Hung Chu² & Irene Shuping Zhao^{1,2}

¹ School of Dentistry, Shenzhen University Health Science Center, Shenzhen, China; ² Faculty of Dentistry, The University of Hong Kong, Pok Fu Lam, Hong Kong, China; ³ Department of Operative Dentistry, RWTH Aachen University, Aachen, Germany

Introduction

Although dentistry has benefitted from technological advancements in recent years, dental caries remains a major oral health problem in most industrialised and non-industrialised countries affecting more than half of schoolchildren and a large majority of adults.¹ Dental caries is a localised chemical dissolution of dental hard tissues caused by the action of acidic by-products of the metabolic processes of cariogenic biofilm.² Conventional non-invasive strategies for caries prevention include oral health education, reduction of sugar consumption, use of fluoride, and application of pit and fissure sealants. Fluoride principally works by hindering the process of demineralisation of enamel and dentine and promoting remineralisation of tooth surfaces with early signs of mineral loss. Pit and fissure sealants generate a physical barrier to prevent the access of dental plaque and its acid products from damaging the enamel surface. However, dental sealants are only effective on the pit and fissures and not the smooth surfaces of teeth. New strategies, such as casein phosphopeptide amorphous calcium phosphate³ and micro- or nanohydroxyapatite compounds, have been proposed to control the balance between demineralisation and remineralisation. It is essential to develop new caries preventive methods to control the disease. A novel, non-invasive approach is the use of laser irradiation on enamel or dentine in preventing caries development. A number of studies reported the potential of laser irradiation on tooth roots or enamel in inhibiting formation of caries lesions.^{4–9}

Several types of laser were studied for caries prevention. The wavelengths of neodymium-doped yttrium aluminium garnet (Nd:YAG) lasers ($\lambda = 1,064\text{ nm}$) and argon lasers ($\lambda = 488\text{--}514\text{ nm}$) make them difficult to be absorbed by enamel. On the other hand, carbon dioxide (CO_2) lasers ($\lambda = 9,000\text{--}11,000\text{ nm}$) are highly absorbed by dental hard tissues and thus have good potential for use in caries prevention.^{7,9} The first CO_2 laser adopted a mixture of nitrogen, helium, and CO_2 , and CO_2 acted

as the active laser medium.¹⁰ Subsequent CO_2 lasers produced various emission laser lines with wavelengths ranging from 9,000 to 11,000 nm. The most common laser lines of CO_2 lasers are centred at 9,300; 9,600; 10,300; and 10,600 nm, respectively. CO_2 laser wavelengths have a higher absorption coefficient to hydroxyapatite than water. A conventional CO_2 laser-emitting light at 10,600 nm is well-absorbed by minerals, while 9,600 nm has the best absorption coefficient to hydroxyapatite followed by 9,300 nm.¹ However, the 10,600 nm CO_2 laser has been commonly used in medicine and dentistry, and most of the commercially available CO_2 lasers operate only at this wavelength. The effect of CO_2 lasers in caries prevention has been studied since the 1960s,¹ when CO_2 lasers (10,600 nm) were discovered to significantly inhibit enamel caries progression.⁷ Significantly less demineralisation was also found in CO_2 laser-treated ($\lambda = 10,600\text{ nm}$) dentine than non-lased dentine.⁹ Furthermore, promising effects of combined CO_2 laser irradiation and fluoride treatment in preventing enamel and dentine caries were reported.^{11–13} However, the mechanism of caries prevention of CO_2 lasers remains to be elucidated. A literature search using the databases PubMed, Scopus and Web of Science revealed no comprehensive review to evaluate studies investigating the effects of actions of CO_2 lasers in caries prevention. Therefore, the objective of this paper was to systematically review the evidence regarding the effects of CO_2 lasers ($\lambda = 10,600\text{ nm}$) on preventing dental caries.

Methods

Search strategy

Two investigators (KL and ISZ) performed a systematic search of articles archived in three databases, PubMed, Scopus and Web of Science. The following keywords were used to identify relevant articles: [(CO_2 laser) OR (carbon dioxide laser)] AND [(dental caries) OR (tooth remineralisation)]. There was no publication year limit, and the last search was made on 31 January 2019. Studies published in English through 2018 and archived in the

PubMed, Scopus, and Web of Science databases were chosen. A list of potentially eligible articles was compiled of publications featuring the keywords (Fig. 1).

Study selection and data extraction

Records identified from database searches were checked for duplication. The titles and abstracts from the potentially eligible list were screened after removing duplicate publications. Articles that did not focus on the effects of CO₂ lasers on preventing dental caries were excluded after screening of titles and abstracts. The remaining articles were retrieved with full texts. The reference lists of all the included articles were screened to identify all possibly eligible studies. The inclusion criterion for selecting studies for this review was that studies investigated the effects of CO₂ lasers on preventing enamel and dentine caries, including their combined effect with topical fluorides on caries progression. For the included papers, the following information was recorded: publication details (authors and years), methods, outcome assessments (various criteria for studying the prevention of caries: reduction of carbonate content, lesion depth, microhardness, mineral loss, surface morphology, and bacterial counts), and the main findings. Any disagreements on study inclusion or data extraction were discussed by the two authors with a third author (OYY) until consensus was reached.

Results

The initial literature search identified 543 potentially relevant articles (143 articles in PubMed, 221 in Scopus, and 179 in Web of Science). A total of 285 duplicate records were removed. After screening the titles and abstracts, 236 articles that were classified as literature reviews, case reports, non-English articles, or irrelevant studies were excluded. No additional relevant publications were found from the references of the selected papers. Finally, 22 articles were found to meet the inclusion criteria and were included in this review. Among these articles, nine studies examined the action of CO₂ lasers on preventing enamel caries (Tab. 1), three studies investigated the effect of CO₂ lasers on preventing dentine caries (Tab. 2), eight studies investigated the effect of CO₂ lasers combined with topical fluorides on enamel (Tab. 3), and three studies examined the effect of CO₂ lasers combined with topical fluorides on dentine (Tab. 4).

Effects of CO₂ lasers on preventing caries of enamel and dentine

Nine studies investigated effects of 10,600nm CO₂ lasers on morphological and chemical changes of enamel as well as on the reduction of enamel demineralisation. Concerning the microhardness analyses, CO₂ laser-treated enamel surfaces showed significantly higher values than those of negative control.^{14,15} Polarised




Fig. 1: Flowchart of the literature search.

light microscopy evaluation revealed that laser-treated enamel groups showed significantly lower lesion depth compared with the non-lased controls,^{16,17} and the decreased lesions represented up to 87% inhibition of the caries progression.⁸ The values of mineral loss of enamel calculated from cross-sectional microhardness analyses were statistically lower in laser-irradiated groups than

Authors, year	Methods	Main findings
Vieira et al. 2015 ¹⁴	Human enamel was irradiated with CO ₂ laser before FTRS and EDX. They were subjected to pH cycling before MHT and SEM.	Irradiated enamel increased microhardness and reduced carbonate content more than negative control. Melting and recrystallisation of enamel occurred after irradiation.
Correa-Afonso et al. 2012 ¹⁵	Human enamel irradiated with CO ₂ laser was subjected to pH cycling before MHT, PLM, and SEM.	Enamel increased microhardness and reduced demineralisation after irradiation. No fusion, melting, or exposure of enamel prisms was found on the irradiated enamel.
Souza-Gabriel et al. 2010 ¹⁸	Human and bovine enamel were irradiated with CO ₂ laser before SEM. They were subjected to pH cycling before MHT.	Enamel showed melting and recrystallisation after irradiation. It was more resistant to demineralisation than the negative control.
Esteves-Oliveira et al. 2009 ¹⁷	Bovine enamel was irradiated with CO ₂ laser before SEM. It was subjected to pH cycling before PLM.	Irradiated enamel showed no ablation, melting, or cracks. It was more resistant to demineralisation than the negative control.
Steiner-Oliveira et al. 2006 ²⁰	Human enamel was irradiated with CO ₂ laser before FTRS and SEM. It was subjected to pH cycling before MHT.	Irradiated enamel had less carbonate content and mineral loss than the negative control. Enamel showed melting and fusion after irradiation.
Klein et al. 2005 ¹⁹	Human enamel was irradiated with CO ₂ laser before SEM. It was subjected to pH cycling before MHT.	Enamel surfaces showed melting and fusion after laser irradiation. They had less demineralisation than the negative control.
Hsu et al. 2000 ¹⁶	Human sound enamel with organic matrix removal or not was treated with CO ₂ laser and subjected to pH cycling before PLM, microradiography, and SEM.	Irradiated enamel had no surface melting or crater after irradiation. It had less mineral loss and lesion depth than other groups. An interaction effect was found between laser irradiation and enamel organic matrix content.
Featherstone et al. 1998 ⁷	Human enamel was irradiated with CO ₂ laser and then subjected to pH cycling before MHT.	Irradiated enamel showed less mineral loss than the negative control.
Kantorowitz et al. 1998 ⁸	Human enamel irradiated with CO ₂ laser was examined by SEM and subjected to pH cycling before MHT.	Irradiated enamel surfaces showed little or no morphological changes. They had less mineral loss than the negative control.

EDX = energy-dispersive X-ray, FTRS = Fourier-transform Raman spectroscopy, MHT = microhardness testing, PLM = polarised light microscopy, SEM = scanning electron microscopy

Table 1: Summary of studies of effects of CO₂ (10,600 nm) lasers on enamel.

those in non-lased groups.^{7,8,16,18–20} There was an interaction effect between the CO₂ laser irradiation and the organic matrix content of enamel on lesion depth and mineral loss of enamel ($p < 0.05$).¹⁶ The enamel with the removal of organic content (non-organic enamel) had a 38% reduction in lesion depth and a 74% reduction in mineral loss after laser irradiation when compared to the non-organic enamel without laser irradiation. Both reductions were significantly lower than the results for the lased normal enamel but greater than those for the non-lased enamel.¹⁶ CO₂ laser irradiation could induce chemical changes in enamel. Fourier-transform Raman spectroscopy showed a reduced carbonate content for laser-treated enamel when compared with non-treated ones,^{14,20} while there were no statistical differences for calcium and phosphorus components between irradiated groups and the control by energy-dispersive X-ray fluorescence spectrometry measurements.¹⁴

Observations under scanning electron microscopy revealed evidence of melting and fusion in the enamel

samples treated with CO₂ lasers,^{14,19,20} and melting and fusion were more frequent in the treated groups that underwent more than one laser application.¹⁴ Enamel surfaces exhibiting melted structures were several times bigger than the prismatic structures and fusion across the prism boundaries.¹⁴ It was verified that a laser-modified layer with a coalescence of crystals was presented in human enamel surfaces, forming an irregular solid mass.¹⁸ A homogeneous and smooth recrystallisation was also observed on the fused enamel surfaces.^{14,18} However, some studies found that the treated enamel areas showed little or even no morphological changes.^{8,16} The irradiated enamel surfaces exhibited a similar appearance to the non-lased controls,¹⁷ and no signs of surface melting, ablation, crater formation, cracks, or fissures could be observed using the parameters showing caries inhibition effects.^{16,17} Three studies investigated the effects of a 10,600nm CO₂ laser on morphological features of dentine as well as on the reduction of dentine demineralisation.^{9,21,22} Root dentine surfaces irradiated with a CO₂ laser with fluences from

Authors, year	Methods	Main findings
de Melo et al. 2014 ²¹	Human root surfaces irradiated with CO ₂ laser were subjected to pH cycling before MHT.	Irradiated root surfaces were more resistant to demineralisation than the negative control.
de Souza-Zaroni et al. 2010 ²²	Human root surfaces were irradiated with CO ₂ laser before SEM. They were subjected to pH cycling before MHT.	Irradiated root surfaces showed melting and resolidification. They showed less mineral loss than negative control.
Nammour et al. 1992 ⁹	Human root surfaces irradiated with CO ₂ laser were subjected to cariogenic challenge before SEM, microdensitometry measurements, and microradiography.	Root surfaces appeared smooth after irradiation. Root surfaces showed melting with no dentinal openings, and they showed more resistant demineralisation than the negative control.

MHT = microhardness testing, SEM = scanning electron microscopy

Table 2: Summary of studies of effects of CO₂ (10,600 nm) lasers on dentine.

4.0 to 6.0 Jcm⁻² showed evidence of melting and resolidification under scanning electron microscopy, and a cracking appearance was more evident on those samples treated with 5.0 and 6.0 Jcm⁻².²² Longitudinally fractured samples had a layer of dentine with no tubular structure (20–70 µm thick), whereas the dentinal tubules retained the normal structure below the sealed layer.⁹ The values of mineral loss were significantly lower in the laser-irradiated dentine groups than the non-irradiated controls.^{21,22} Additionally, laser-induced inhibitory

effects on dentine demineralisation were observed when fluences reached or exceeded 4.0 Jcm⁻².

Effects of CO₂ lasers combined with topical fluorides on enamel and dentine

Eight studies investigated the combined effects of 10,600 nm CO₂ laser irradiation and the application of topical fluoride on enamel. The results of these studies varied. Four studies found that there was a synergistic

AD

SUBSCRIBE NOW!

ceramic implants—international magazine of ceramic implant technology

Interdisciplinary, transparent & market orientated



ONLINE
SUBSCRIPTION HERE



www.oemus-shop.de

Fax: **+49 341 48474-290**

I would like to subscribe for the following journals:

- ☐ ceramic implants—international magazine of ceramic implant technology 2 issues p.a. € 30*
- ☐ implants—international magazine of oral implantology 4 issues p.a. € 44*

Terms & Conditions: The subscription may be cancelled in written form without due justification within 14 days of order by contacting OEMUS MEDIA AG, Holbeinstraße 29, 04229 Leipzig, Germany. Dispatching notification in good time will suffice. The subscription is automatically extended by another twelve months if it is not cancelled in written form 6 weeks prior to the end of the reference period.

* All prices include VAT, shipping and handling.

Last Name, First Name

Company

Street, ZIP, City, Country

E-mail

Credit Card Number

Expiration Date

Security Code

Stamp

Signature

Holbeinstraße 29 · 04229 Leipzig · Germany · Phone: +49 341 48474-201 · s.schmehl@oemus-media.de